

# Abortion Laws and Women's Health\*

Damian Clarke<sup>†</sup>

Hanna Mühlrad<sup>‡</sup>

September 24, 2020

## Abstract

We examine the impact of progressive and regressive abortion legislation on women's health in Mexico. Following a 2007 reform in the Federal District of Mexico which decriminalised and subsidised early-term elective abortion, multiple other Mexican states increased sanctions on illegal abortion. We observe that the original legalisation resulted in a sharp decline in maternal morbidity, particularly morbidity due to haemorrhage early in pregnancy. We observe small or null impacts on women's health from increasing sanctions on illegal abortion. These results quantify the considerable improvements in non-mortal health outcomes flowing from legal access to abortion.

**JEL Codes:** I18; J13; K38; H75.

**Keywords:** Abortion; Maternal Morbidity; Maternal Mortality; Health Care Provision; Political Economy; Legislative Reform.

---

\*We thank the Editor Ana Balsa, two anonymous referees, Sonia Bhalotra, René Castro, Blair G. Darney, Dolores de la Mata, Hans Grönqvist, Randi Hjalmarsson, Lakshmi Iyer, Elin Larsson, Jason Lindo, Andreea Mitrut, Carol Propper and seminar audiences at IPAS Mexico, The LACEA Health Economics Network, the IZA Gender Workshop, CIDE Mexico, Colegio de Mexico, the Mexican Institute of Public Health, University of Gothenburg, SOFI Stockholm University, Karolinska Institute and CSAE Oxford. We thank Raffaella Schiavon for detailed discussions regarding the practical implementation of Mexico D.F.'s ILE reform and her generosity in sharing expertise. We are also grateful to Natalia Volkow at INEGI for providing access to disaggregated hospitalisation data, Alejandro del Valle for sharing data, and Cristhian Molina for excellent research assistance. Clarke acknowledges the generous support of FONDECYT Regular, grant number 1200634 from the Government of Chile and the institutional support of the Millennium Institute for Market Imperfections and Public Policy. This paper considerably extends and partially subsumes an earlier study by these authors called "The Impact of Abortion Legalization on Fertility and Female Empowerment: New Evidence from Mexico".

<sup>†</sup>Department of Economics, Universidad de Chile and IZA. Diagonal Paraguay 257, Santiago, Chile. Contact: dclarke@fen.uchile.cl.

<sup>‡</sup>PhD, Department of Clinical Sciences, Danderyd Hospital (KI DS) | Karolinska Institutet, Sweden. Contact: hanna.muhlrad@ki.se.

# 1 Introduction

Appeals to women's health are frequently made when debating the merits of abortion legislation. These calls are made by both advocates of legal abortion, as well as those advocating for abortion to become, or remain, illegal. The arguments backing up such claims are drawn from a range of sources, which are often correlational or based on small or non-representative samples of women.<sup>1</sup>

In this study, we present the first population-level evidence of the impact of sub-national variation in abortion laws on maternal *morbidity*, as well as mortality, using the universe of administrative health records from Mexico.<sup>2</sup> We focus on a period in which considerable within-country reform of abortion policy was undertaken, with both a sweeping legalisation in the Federal District of the country (Mexico DF), and increasing sanctions on (already illegal) abortion in other regions. In this context, we are able to determine to what extent change in abortion laws, absent other major contraceptive revolutions, impact health indicators for the population of affected women. We combine the state-level variation over time resulting from legislative changes in abortion law with high-quality vital-statistics data recording over 30 million births, 18.4 thousand maternal deaths and 46 million inpatient cases for causes related to maternal health. To seek to understand the observed impact of abortion laws on health outcomes, we also take advantage of rich administrative and survey data to identify the reform's wider impact on birth rates, judicial sentencing, and on sexual behaviour, and explore potential channels of health impacts, including improvements in the quality of abortion care, and changes in the composition of women giving birth.

The environment under study provides a unique opportunity to examine simultaneous expansions and contractions of abortion policies.<sup>3</sup> While much of the existing literature on the impact of

---

<sup>1</sup>The use of such arguments even when based on weak evidence is not isolated to non-governmental organisations. Similar arguments are also made by politicians. One such example is a fact sheet published on the US National Cancer Institute website by the Bush administration positing an (unfounded) link between abortion and breast cancer (Special Investigations Division, Committee of Government Reform, House of Representatives, 2003).

<sup>2</sup>Associations between abortion legalization and maternal mortality or morbidity have been documented in the medical and public health literature for multiple countries (Benson et al., 2011) including Albania (Sahatci, 1993), Bangladesh (Chowdhury et al., 2007), Nepal (Henderson et al., 2013), Romania (Serbanescu et al., 1995; Stephenson et al., 1992), Singapore (Singh and Ratnam, 2015) and South Africa (Rees et al., 1997). Existing studies are mainly based on reviews of medical charts at selected hospitals, for example in the US (Goldstein and Stewart, 1972; Stewart and Goldstein, 1971; Seward et al., 1973; Kahan et al., 1975), Guyana (Nunes and Delph, 1997), Nepal (Henderson et al., 2013) and South Africa (Mbele et al., 2006; Jewkes et al., 2002). We are aware of no prior studies which are based on population-level data, and based on within-country variation in abortion reforms.

<sup>3</sup>As we discuss at more length in Sections 2.2-2.3 of this paper, the change due to the initial legalisation of abortion in Mexico DF was considerably larger than subsequent legislative tightenings in other Mexican states. In the case

abortion—and contraceptive policies more generally—focuses on expansions in access, there are a number of papers which focus on contractions in policies. These include historical restrictions in Romania (Pop-Eleches, 2010), the impact of parental consent or notification laws targeted at adolescents in the U.S. (Bitler and Zavodny, 2001; Joyce and Kaestner, 1996), and a recent hollowing out in the availability of providers due to state-specific legislation in the U.S. (Fischer et al., 2018; Lindo et al., 2019). However, the legalisation of abortion in Mexico DF, and the resulting spate of constitutional changes increasing the harshness of sentencing of illegal abortion, provides the opportunity to examine the impact of a contemporaneous series of restrictive and permissive abortion policies in a single country and time.

Reproductive health rights have been documented to be of considerable economic significance. This fact has been emphasized by a growing body of economic literature which has—empirically and theoretically—demonstrated how access to elective abortion and contraceptive methods has shaped fertility patterns, marital markets, crime, education, the labor market and female empowerment (e.g. Ananat et al., 2007; Bailey and Lindo, 2017; Chiappori and Orefice, 2008; Guldi, 2008; Mitrut and Wolff, 2011; Myers, 2017; Pop-Eleches, 2010). The impact of abortion laws on women’s health has received less attention, and the causal relationship is yet to be established. To the best of our knowledge, this is the first study to provide well-identified population-level evidence of the impact of abortion legalisation on maternal morbidity and mortality based on within-country variation in abortion availability. While an association between abortion legalization and lower abortion-related complications has been documented in previous studies, comprehensively capturing the impact of the passage of abortion law on abortion-related morbidity is a considerable challenge, especially in clandestine settings, where under-reporting may occur (Singh et al., 2010).<sup>4</sup>

Using two-way fixed effect (FE) models and panel event study methods (and synthetic controls as a robustness check), as well as recent advances in a literature examining causal estimation with time varying policy reform (Rambachan and Roth, 2019; Goodman-Bacon, 2018), we observe consistent evidence to suggest the Mexico DF’s abortion legalization brought about a sharp reduction in fertility

---

of the constitutional changes issued by states, in each case abortion was already illegal, and any changes owe to an increased threat of prosecution or sanction. Using the universe of legal decisions in the country, we do document evidence suggesting that these reforms increase the average length of sentences handed down to women.

<sup>4</sup>Maternal mortality is considered the “tip of the iceberg”, where the mass consists of maternal morbidity (Loudon, 1992). In many settings, analyses of the impact of abortion on population health focuses only on maternal mortality due to a lack of universal health records measuring maternal morbidity.

(by 8%), haemorrhage early in pregnancy (by 35%) and abortion related morbidity (by 20%). Event study estimates examining increases in sanctions on (already illegal) abortion point to much smaller effects on these variables, which are generally not statistically significant. In general, we observe impacts on maternal *mortality* which are hard to consistently sign, given that they are considerably less precise, suggesting that when only examining impacts of abortion law on maternal mortality, analyses fail to account for the full weight of abortion reform on women's health.

Previous studies on Mexico's abortion reform laws include legal and ethical overviews (Johnson, 2013; Madrazo, 2009), qualitative studies on abortion provision (Contreras et al., 2011; Schiavon et al., 2010), quantitative studies on abortion services and patient characteristics using data from selected hospitals or surveys (Mondragón y Kalb et al., 2011; Becker et al., 2013), and fertility trends using vital statistics (Gutierrez-Vazquez and Parrado, 2015).<sup>5</sup> This paper contributes to previous studies by examining the causal impacts of abortion legalization, as well as regressive law changes, on women's health and well-being using the full power of vital statistics data including administrative microdata on births, maternal morbidity, maternal mortality and judicial statistics on penal matters. In addition, we examine heterogeneous effects of the reform as well as potential mechanisms such as usage and knowledge of contraceptive methods.

There is a large unmet need for family planning in low- and middle-income countries (Sully et al., 2019). Yet, the evidence on the impacts of reproductive health rights including safe and legal abortions in these settings, especially building on microdata, is very scarce. In light of this, our study provides strong evidence that abortion legalization in an emerging economy leads to rapid and discernible changes in political behavior, aggregate fertility rates, and (significant improvements in) maternal health. These results provide a number of important policy lessons for jurisdictions considering changes in abortion laws.

---

<sup>5</sup>In examining the abortion reform and fertility outcomes, Gutierrez-Vazquez and Parrado (2015) use national vital statistics to examine the effect on fertility across ages. Due to the use of a limited amount of data and limitations inherent in the empirical design one cannot assign a causal interpretation to the results with confidence. More specifically, only three different years of data are used (1990, 2000 and 2010). In a study by Koch et al. (2015), maternal mortality is found to increase in areas with more liberal abortion laws. Koch et al. (2015) however, has received strong criticism for highly misleading and inaccurate data selection (Darney et al., 2017).

## **2 Background**

### **2.1 Unsafe Abortions and Maternal Mortality in Mexico**

Globally, maternal mortality has declined from 385 maternal deaths per 100,000 live births in 1990 to 211 deaths per 100,000 live births in 2017. The overwhelming majority of these deaths occurred in low income countries. During the same period in Mexico, there has been a similar decline in maternal mortality from 88 to 33 deaths per 100,000 live births (Rodríguez-Aguilar, 2018; Bongaarts, 2016). However, Mexico has still not achieved its 2015 Millennium Development Goal, of a ratio below 22 deaths per 100,000 live births. In Mexico, the highest prevalence of maternal mortality is found in metropolitan areas and among women aged 20-34. Rates of maternal mortality is especially high in socially vulnerable populations across Mexico. Areas with the highest rates of maternal mortality exhibit some of the lowest levels of government expenditure on maternal health (Pérez-Pérez et al., 2019).

Abortion-related mortality represents a substantial proportion of all maternal deaths in Mexico, accounting for 7.2 % of all maternal deaths during 2000-2008 (Schiavon et al., 2012a). While there has been a downward trend in maternal mortality in Mexico between 1990-2008, maternal mortality attributed to abortion-related causes has not exhibited similar declines during this period. The majority of these deaths occur in women without health insurance (Schiavon et al., 2012a).

The rate of abortion-related mortality varies substantially across regions of Mexico. The highest rates can be found in some of the poorest states, such as Guerrero and Chiapas with as many deaths as 140 and 90 per 100,000 hospitalizations. This can be compared to Baja California Sur with only 9 deaths per 100 000 hospitalization. Yet, high rates of abortion-related mortality are not exclusively observed in poor states across Mexico. In Mexico City, for example, the abortion-related mortality rate was 38 per 100,000 hospitalizations during 2000-2008.

Unsafe abortion procedures account for most abortion-related mortality (Schiavon et al., 2012a). Rates of unsafe abortion are particularly high in the Latin America and Caribbean region with an estimated 4.2 million unsafe induced abortions being performed each year (World Health Organization, 2011). This region also exhibits some of the world's most conservative laws on abortion (United Nations, 2014). However, restrictive laws on abortion do not translate to lower rates of induced abortion but are instead associated with higher rates of unsafe abortion and correspond-

ing higher rates of abortion-related morbidity and mortality compared to settings with more liberal access policies (Guttmacher Institute, 2012).

Indeed, the rate of induced abortion in Mexico is considered high internationally (Becker, 2013). Despite high rates of contraceptive use, the estimated rate of induced abortions increased from 1990, with 25 abortions per 1,000 fertile-aged women, to 2006, with 33 abortions per 1,000 women (Juarez et al., 2008). Many of these induced abortions are performed in clandestine and often unsafe settings (Guttmacher Institute, 2012). Medical records from public hospitals in Mexico show high rates of abortion-related complications, with an estimated 150,000 women treated for complications in 2006 alone (Juarez et al., 2008).

## **2.2 The 2007 legal interruption of pregnancy reform in Mexico DF**

Since the 1970s, women's rights advocates in Mexico have been promoting women's health rights including access to safe abortions (Kulczycki, 2011). To address the issue of unsafe abortions as a preventable cause of maternal morbidity and its huge burden on public health system, a National Pro-choice Alliance in Mexico was established in 2000 to promote women's sexual and reproductive health rights. This movement was supported by a wide range of groups including health care professionals, women's rights groups and activists, politicians, academics and Catholics with pro-choice views (Johnson, 2013; Madrazo, 2009; Kulczycki, 2011; Blanco-Mancilla, 2011).<sup>6</sup>

With the support of the leftist PRD party, the legislative assembly of the Federal District of Mexico voted to legalize elective abortion (termed legal interruption of pregnancy, or ILE for its name in Spanish) on April 24, 2007 (Kulczycki, 2011). The ILE reform was signed into law the following day, and published in the official Gazette of the Federal District on April 26, 2007 (Ciudad de México, 2007). This immediately permitted women above the age of 18 to request an abortion at up to 12 weeks of gestation without restriction and free of charge. Access for minors requires parental or guardian consent. Under this law, induced abortion was made legal in both the public and private health care sectors (though requiring payment in private clinics).

---

<sup>6</sup>This alliance was successful in moving the women's health agenda rapidly forward in the historically conservative setting on Mexico City using a number of different strategies including the involvement of health care officials, hospitals and other health care providers, which was not only important for building support for the reform, but also to facilitate a smooth implementation of the abortion program. They were also successful in gaining a strong presence in the public media for promoting women's health rights (Kulczycki, 2011; Johnson, 2013).

Immediate implementation was made possible by collaboration between the Ministry of Health of Mexico DF, members of the health department and international NGOs, which had thoroughly designed a program for public provision of abortion services called the “the ILE program” and its implementation even before the law was passed (Singh et al., 2012a). As such, abortion services were made available via public clinics immediately after the law was passed in April 2007, although with lower capacity and efficiency compared to current conditions. Abortion services were also quickly available in the private health care sector (Blanco-Mancilla, 2011). Additionally, under this law sexual education in schools was improved, and post-abortion contraceptives were made freely available directly from the health clinics which provided abortions (Contreras et al., 2011). On August 29, 2008 the decision to pass the ILE law was ratified by the Supreme Court of Mexico, making Mexico DF, together with Cuba and Uruguay, the most liberal jurisdiction in terms of abortion legislation in the entire Latin American and Caribbean region (Fraser, 2015).

Any abortions conducted in publicly run clinics are provided free of charge to residents of Mexico DF. Women with residency outside Mexico DF can also access the public provision of abortion through the Ministry of Health in Mexico DF (MOH-DF) but are charged based on a sliding fee scale depending on the woman’s socioeconomic background. In 2010, 74% of all women who received an abortion through the public health care sector were women living in Mexico DF, 24% were living in the state of Mexico (which shares a border with Mexico DF) and 2% were living in other states (Mondragón y Kalb et al., 2011). The age profile of women seeking abortions largely mirrors the age-profile of births, occurring at slightly greater rates among younger (under 25 years) and older (above 36 years) groups (Appendix Figure A1).

Records from public hospitals show that during the year of 2007, when the reform was implemented, more than 7,000 abortion procedures were performed at 14 selected MOH-DF clinics. Over the years, the MOH-DF abortion program expanded its services and became more efficient in meeting the high demand for elective abortion. The MOH-DF program offers both surgical and medical abortion procedures and is the main provider of medical abortion (Winikoff and Sheldon, 2012). The large shift from 25% of all abortion procedures being medical in 2007 to as much as 74% in 2011 have played a key part of meeting the demand and reducing complications and side-effects (Becker,

2013).<sup>7</sup> As of 2015, approximately 150,000 abortions were carried out at the MOH-DF clinics.<sup>8</sup> Information regarding the private provision of abortion services is limited due to a lack of supervision of the private market for legal abortion services (Becker, 2013). Despite the fact that safe abortion, at no or low cost, is provided by the public health system in Mexico DF, women do seek abortion services within the private sector. A descriptive study by Schiavon et al. (2012b) suggests that private abortion services are provided at high costs (157–505 US dollars) and quality of care is inferior to that in the public sector, given that the less safe and efficient “dilation and curettage” is used as the main method in the private sector (71%).

### **2.3 Regressive law changes following the ILE Reform**

Almost immediately following Mexico DF’s ILE reform, a number of states began a series of counter-legislations to change the respective sections of their constitutions or penal codes, defining the beginning of human life as occurring at conception. Often, these legal responses directly referenced Mexico DF’s ILE reform.<sup>9</sup> Even in cases where they did not directly refer to the ILE reform, it seems highly likely that the reform was a defining factor. For example, in the 20 years prior to the ILE reform there had been only two constitutionally defined changes to the articles relating to abortion in the penal codes of all states of Mexico (Gamboa Montejano and Valdés Robledo, 2014), compared to 18 changes between June 21, 2008 and November 17, 2009. Importantly, these reforms resulted in constitutional changes which recognised life as beginning at conception, opening the door for potential homicide charges. In order to understand the precise nature of law changes, we conducted a side-by-side reading of penal codes from pre- and post-reform for each state undertaking an abortion reform. The nature of changes implied by the reform are documented in Appendix Table A1. As we display there, of the states undertaking a regressive reform, all but 6 formally altered their penal codes to change sanctions in the case of proven abortions, while the remaining 6 only altered their state constitution to recognise life as beginning at conception.

---

<sup>7</sup>Misoprostol alone was the main regimen for medical abortions in MOH-DF until 2011 when Mifepristone (combined with Misoprostol) was introduced, making the medical abortion procedures provided by the ILE program more efficient and safe.

<sup>8</sup>To put this in context, we note that the quantity of abortions per year (adjusted for population) is similar to the quantity in the United States. In 2010 for example, 16,945 abortions were provided by the ILE program. In 2010 Mexico DF had approximately 8.55 million inhabitants. Thus, adjusted to the US population in 2010 (308.7 million), this would imply 611,803 abortions. In reality, 765,651 abortions were reported in the US in 2010 (Pazol et al., 2014).

<sup>9</sup>For example, the constitutional decree issued by the state of Nayarit when changing their penal code explicitly refers to the changes in the penal and civil code of Mexico DF (Gobierno de Nayarit, 2009, p. 14).



In Appendix Figure A2, we display the geographical distribution of law changes (progressive, regressive or neutral) over the period under study. The only progressive reform refers to Mexico DF’s ILE reform, while 18 states made regressive changes (ie legal tightenings) after the initial reform. We have compiled on a state-by-state basis the exact dates the reforms were passed into law, and these are displayed in Appendix Table A2. To the best of our knowledge, there exists no centralized record of the dates and laws which were altered in the post ILE era, and as such we compiled these from our reading of legal source documents. In Section 4 of this paper we return to how we use the geographic and temporal variation in the passage of laws in our identification strategy.

## **2.4 The Potential Channels for Impacts of Abortion Reform**

Legislative reforms to abortion policies could potentially impact health through a number of channels. We lay these out conceptually below, without yet seeking to assess their relative importance or relevance in explaining observed outcomes. We then move on to assess the likelihood that particular channels can explain observed impacts of abortion reform on maternal health (in the context of this paper) in Section 5.3.

**1. Unsafe abortions are shifted to safe legal conditions** This channel is relevant for individuals who would abort when abortion is illegal, and also abort when abortion is legal. As discussed in Section 2.1 unsafe abortion is thought to be a significant determinant of maternal death and hospitalisation, with evidence that a lack of access to safe abortion shifts the demand for abortion into clandestine and unsafe conditions (Haddad and Nour, 2009). To the degree that legalizing abortion shifts clandestine abortions at risk of complications to sanitary conditions with lower risk, there exists a hypothesised direct channel through which abortion reform may impact women’s health. In this channel, holding all else constant, abortions which would have occurred whether abortion was legal or not should imply a direct improvement in women’s health when abortion is legalised. We refer to this as the “quality of care channel”.

**2. Undesired pregnancies are avoided** This channel is relevant for individuals who would not abort when abortion is illegal, but abort when abortion is legalized. To the degree that abortion being illegal discourages women who would otherwise have sought to discontinue their pregnancy

from seeking abortion, changes in laws will shift the demand for abortion. This channel, which we refer to as the “demand” channel could, theoretically at least, have varying impacts on women’s health. These are (a) for undesired pregnancies which would have resulted in poor maternal health outcomes, the legalization of abortion and resulting avoidance of risky pregnancies will, all else constant, improve maternal health. However (b) for undesired pregnancies which would not have resulted in poor maternal health outcomes, the legalization of abortion may shift maternal health outcomes if abortion – even in safe legal conditions – implies some risk of complications.<sup>10</sup>

**3. The composition of women becoming pregnant changes** This channel is relevant for individuals who would not abort when abortion is illegal, and also not abort when abortion is legal. This channel is relevant only to the degree that the population of women giving birth following abortion reform will consist of more women who desire a birth, regardless of whether abortion is legal. As women’s health complications arise throughout pregnancy, changes in the composition of women becoming pregnant and taking a pregnancy to term may result in changes in rates of health complications observed at the population level. Ex ante, this selection process has ambiguous impacts on maternal health, as it depends on the nature the selectivity. We will refer to this as the “selection channel”.

### 3 Data

We construct a balanced panel recording morbidity and mortality outcomes, birth rates and legal outcomes for each of Mexico’s 32 states between the years of 2001-2015. In principal models we consider a monthly panel, however also generate more temporally aggregate data (by year and by trimester) to include the universe of cases where monthly records are not available, as laid out below. We construct this from a large number of administrative microdata bases covering 100s of millions of records, which are described at more length in Appendix B and summarised below. Along with outcomes generated from administrative data, we collect a number of measures of sexual behaviour from survey data, time varying controls, and the exposure to the ILE reform or subsequent legal changes.

---

<sup>10</sup>As we discuss later in the paper, this second channel appears to be quite unlikely given that abortion in safe settings has very low rates of associated morbidity and mortality. In fact, induced abortion in safe setting are considered “one of the safest procedures in contemporary practice” with a mortality rate below 1 per 100,000 procedures (Grimes, 2005).

**Health Outcomes** We observe all morbidity events resulting in a hospital inpatient visit to any public hospital in Mexico. Depending on the hospital type (state run or social-security run), the microdata format varies, and as such, we can consistently group records only for the years 2004-2015, and only with a yearly total. Given this, we provide two sets of results: one based only on visits to state-run hospitals, as in this case we can calculate monthly aggregates for the longer period of 2001-2015, and another based on yearly records for the shorter 2004-2015 period when all hospitalisation data is observed. We similarly consider leakage to the private system using administrative records from all private hospitals.

Our principal measures of interest are maternal morbidity outcomes, classified according to International Classification of Diseases tenth revision (ICD-10) codes,<sup>11</sup> plausibly impacted by abortion. Specifically, these are (a) “abortion related morbidity” (ICD-10 codes O02-O08), and (b) haemorrhage early in pregnancy (ICD code O20).<sup>12</sup> The remainder of the ICD codes are not examined as it is unlikely that they are sequelae of abortion (for example eclampsia or pre-eclampsia), or are morbidities occurring in the puerperium period, and so unable to be sequelae of abortion. These two codes are not chosen arbitrarily by the authors, but rather in line with the fact that haemorrhage and incomplete abortion are the two most common complications of unsafe abortion (World Health Organization, 2018). We provide a full break-down of all ICD codes related to pregnancy, childbirth and the puerperium (the ICD-10 O codes) in Appendix Table A3 which documents both the considerable frequency of the two chosen classes, as well as the implausibility of impacts on other variables. Finally, as an exploratory analysis we aim to measure mental health outcomes of women. In practice, the only way that we can consistently measure this in our data is via the code F53 which captures post-partum depression.<sup>13</sup>

---

<sup>11</sup>It is important to note that the procedure of assigning ICD codes for a hospitalization is typically implemented by a particular person or persons in charge of coding diagnoses in hospitals (such as an epidemiologist or a hospital employee). This coding is made based on their reading of the treating team’s charts, and is *not* determined by the team actually treating the patient (Velasco Sustaita, Undated). Moreover, the ICD coding is typically processed after the patient is discharged from the hospital. Anecdotal evidence suggests that the prosecution of women for the crime of abortion usually occurs during the hospital stay, after a clinician/nurse/social worker have raised “suspicion” to the police. There is no ICD-code of induced illegal abortion in the ICD-10 classifications, and indeed, evidence from other contexts suggests that if anything, providers may seek to record abortion related morbidity within sub-codes of ICD-10 classifications (such as recording abortions as spontaneous instead of induced), rather than classifying as an alternative code (Suh, 2014). The authors are grateful to Dr. Raffaella Schiavon for providing very useful background details in correspondence.

<sup>12</sup>For a longer discussion on classification of abortion-related morbidity see Schiavon et al. (2012a) and World Health Organization (2018).

<sup>13</sup>We additionally explored using external causes of morbidity to capture suicide attempts, however the inpatient

We additionally consider a number of placebo outcomes. We do this in a number of ways. A first placebo test consists of considering a (presumably) unaffected morbidity cause *within* the same ICD class (the O codes), namely late term obstetric complications (ICD codes O70-O75). While this is not a perfect placebo insofar as it may be impacted by changing composition of mothers (via the ‘selection’ channel indicated earlier), it is unlikely to be mechanically related to abortion, so provides a useful comparison to our main morbidity outcomes. Additionally, and to completely isolate the placebo tests from potential changes in composition of mothers, we consider a larger number of placebo outcomes coming from *other* ICD classifications.<sup>14</sup>

While our main interest here is to document the impacts of abortion reform on the much larger pool of morbidity outcomes, we additionally consider maternal *mortality* as recorded in Mexico’s complete vital statistics registers. We thus calculate the number of all maternal deaths in each state and year, and additionally only those maternal deaths classified as owing to abortion. Additional discussion of the generation of these data, as well as the recording of microdata bases is provided in Appendix B.

**Births** In order to benchmark the Mexican abortion reforms’ impact on birth rates with respect to the wider literature, we also require aggregate data on birth rates by state. We generate these state-level measures from publicly available microdata on births provided by INEGI. We use each birth register occurring to women aged 15–49 over the time period of 2001–2013; a sample of 30,341,376 births. As discussed at more length in the Data Appendix, we can only observe birth records up to the year of 2013 as we follow the procedure suggested by the National Population Council (CONAPO) of using birth registers up to four years following each date to ensure that we capture births even if they are registered with a considerable delay. We use these same data to measure average characteristics of mothers and fathers (where registered) to examine any changes in composition of parents following abortion reforms. In both the case of births and the case of health outcomes, in order to calculate rates of occurrence we record state.level population in each year for all women aged 15–49 provided

---

data from Mexico only consistently provides a single ICD code capturing the principal cause of hospitalization, while external causes are classified as a secondary ICD code. Thus, these external causes are not recorded consistently in administrative data.

<sup>14</sup>Specifically, these are Diseases of the ear and mastoid processes; Neoplasms; Endocrine, nutritional and metabolic diseases; Diseases of the blood and bloodforming organs; Diseases of the nervous system; and Diseases of the skin and subcutaneous tissue. We discuss these at more length in Section 5.3.

by CONAPO.

**Additional Measures** We additionally collect a number of other state- and time-varying measures to examine potential mechanisms of action of the reform, or to dismiss alternative possible explanations of observed impacts. These measures consist of: (a) all first legal decisions made by the Mexican judiciary related to abortion compiled from microdata collected by Mexico’s Judicial Statistics on Penal Matters; (b) information on contraceptive use and sexual behaviour from the Mexican Family Life Survey (MxFLS); and (c) time-varying controls to capture possible confounders of abortion policy, namely education, health investment and access including access to the public insurance program *Seguro Popular*, economic development, and women’s social inclusion. These controls are fully described in Data Appendix B, as well in discussion of summary statistics below.

**Summary Statistics** Summary statistics of principal outcomes (maternal morbidity, mortality and births) are provided in Table 1. The total number of cases of each morbidity class are described in panel A, and mortality outcomes, both for all maternal deaths, and those only classified as owing to abortion, are provided in panel B. State by year×month averages of the number of births and births per 1,000 fertile-aged women are displayed in panel C of Table 1. On average, morbidity outcomes are various orders of magnitude higher than mortality outcomes. For example, the average quantity of hospitalisations for abortion related causes was 269 per state and month, versus 3 maternal deaths on average, or 0.2 maternal deaths when considering only those classified as owing to abortive causes. In this Table, morbidity data is recorded only based on public hospital data where the month of the visit is recorded. As we will discuss at more length below, we will also consider variation by state and year allowing us to capture the full universe of health outcomes from both state-run and social security run hospitals (see Appendix Table A4 for summary statistics), and control variables are summarised in Appendix Table A5.

Raw trends of principal maternal health outcomes are provided in Figure 1, pointing to important shifts in inpatient visits related to maternal health following the ILE reform. Figure 1(a) and (b) document the quantity of monthly inpatient cases for abortion related morbidity and haemorrhage in early pregnancy in Mexico DF, states that adopted a subsequent legal tightening, and states with no subsequent legal changes. We observe reductions in the absolute quantity of cases in Mexico

DF in both cases following the ILE reform in April 2007 (indicated as a vertical red line). Panels (a) and (b) are based only on hospitalization data from Secretary of Health hospitals. When we additionally extend to include Social Security hospitals in panels (c)-(f), observing yearly variation only, we see a similar pattern with reductions in total hospitalisations (panels (c) and (d)), and rates of hospitalisations for fertile-aged women (panels (e) and (f)).

Table 1: Summary Statistics for Month by Year by State Specifications

Variable	Obs.	Mean	Std. Dev.	Min.	Max.
<b>Panel A: Morbidity Outcomes</b>					
Total Number of Deliveries in Public Hospitals	5760	1455.9	1282.8	143	8496
Total Inpatient Cases for ICD O codes, except births	5760	1276.3	1002.8	139	6271
Total Inpatient Cases for Abortion-Related Causes	5760	268.7	232.9	27	1573
Total Inpatient Cases for Haemorrhage Early in Pregnancy	5760	31.4	27.2	0	207
Total Inpatient Days for Abortion-Related Causes	5760	361.3	327.4	28	2168
Total Inpatient Days for Haemorrhage Early in Pregnancy	5760	64.7	63.0	0	463
Total Inpatient Cases for Obstetric Complications	5760	36.0	51.1	0	375
Total Inpatient Cases for Post-Partum Depression	5760	0.1	0.3	0	6
<b>Panel B: Mortality Outcomes</b>					
Total Number of Maternal Deaths	6144	2.98	3.26	0.00	25.00
Total Number of Maternal Deaths due to Abortion	6144	0.21	0.50	0.00	4.00
<b>Panel C: Demographic Outcomes</b>					
Population of 15-49 Year-old Women	6144	864691	743706	116430	4228223
Total Number of Births	4992	6078	4904	719	28546
Birth rate per 1,000 women	4992	7	1	5	13

Each observation is a state×year×month cell. Mexico is composed of 32 States. The number of cells varies due to the number of years and months of data available. In panel A, morbidity data is displayed for 12 months in 12 years (2004-2015). Values are generated from all inpatient cases as classified from microdata from the primary care (hospital) records from all public hospitals administered by the Secretariat of Health (Social Security System hospitals do not report month of hospitalization). Each type of morbidity is classified by ICD-10 codes. In Panel B, mortality outcomes are displayed for 12 months in 16 years (2001-2016). In panel C, data on population is displayed for 12 months in 16 years (2001-2016), and data on births is displayed for 12 months in 13 years (2001-2013). Following CONAPO, the last four years of birth outcomes are suppressed to account for reporting outside of the period of birth. State×year summary statistics including Social Security System hospitals are provided in Appendix Table A4.

## 4 Methodology


We aim to examine the impact of the ILE reform and regressive law changes, compared with outcomes in states in which no reform was implemented. We thus begin by estimating the following two “two way fixed effect” specifications<sup>15</sup>:

$$Health_{st} = \alpha_0 + \alpha_1 ILE_{st} + X'_{st} \Gamma_1 + \phi_s + \mu_t + (\phi_s \times month) + \varepsilon_{st} \quad (1)$$

$$Health_{st} = \beta_0 + \beta_1 Regressive_{st} + X'_{st} \Gamma_2 + \phi_s + \mu_t + (\phi_s \times month) + \eta_{st} \quad (2)$$

<sup>15</sup>We estimate these models separately, however all results documented in this paper are robust to estimating a single model including both the ILE and Regressive variables. Full results in this set-up are documented in Clarke and Mühlrad (2018).

Figure 1: Raw Trends in Abortion Related Morbidity and Haemorrhage Early in Pregnancy



Notes: Figures present the total number of discharges due to abortion related morbidity (left-hand panels) and haemorrhage early in pregnancy (right-hand panels) in Mexico DF (which adopted the ILE reform), states which had a later tightening of abortion laws, and all other states. Panels (a) and (b) plot monthly trends based on all discharges from Health Secretariat hospitals given that these are the only registers which record month and year of discharge. Panels (c) and (d) plot yearly trends based on all public hospital discharges (both Health Secretariat hospitals and Social Security Hospitals which do not report month of discharge), and panels (e) and (f) plot yearly discharges based on all public hospital data expressed per 1,000 women of fertile age. The dotted vertical line is plotted in April of 2007, the date of passage of the ILE abortion reform, and wide-scale rollout of available abortions. Residualized plots and longer yearly plots based on panels (a) and (b) are provided as Appendix Figures A3 and A4 respectively.

Here *Health* refers to average rates of morbidity or mortality in state  $s$  at time  $t$ , and ILE and Regressive refer to the post-ILE and post-Regressive Law changes in affected states. Our parameters of interest are  $\alpha_1$  from equation 1 and  $\beta_1$  from equation 2. In the case of specification 1, the estimation sample consists of Mexico DF (which adopts ILE) and all non-reforming states, and in the case of specification 2, the sample consists of all states which adopt regressive abortion laws and all non-reforming states. We include state and year fixed effects as  $\phi_s$  and  $\mu_t$  respectively as well as state-specific monthly fixed effects ( $\phi_s \times month$ ) to capture any potential state-specific seasonality, and examine stability to the inclusion of the time-varying controls  $X_{st}$  listed in Section B.2. In initial specifications, time  $t$  refers to monthly measures. This allows us to examine fine-grained temporal variation in outcomes in all hospitals run by the Mexican Secretariat of Health. However, to ensure that these results are also observed in the full universe of public health records, we also consider specifications where time  $t$  refers to yearly cells including all data.

In this two-way FE specification, the ILE variable in equation 1 is an indicator which switches from 0 to 1 at a particular point in time, however the Regressive indicator in equation 2 switches from 0 to 1 at varying times depending on the state. In the case of heterogeneous treatment effects and time-varying adoption, this single-coefficient model can considerably mis-estimate the average treatment effect on the treated (ATT), given that already treated units act as controls in future periods (Goodman-Bacon, 2018). While this is not of concern for our estimate of  $\alpha_1$ , it is for our estimate of  $\beta_1$ , and so we will: (a) document the parameter decomposition derived in Goodman-Bacon (2018), and (b) additionally estimate event study specifications which avoid these potential biases. There are 32 states in Mexico (including the Federal District), and these laws are defined at the level of the state. In order to account for the possibility of unobserved correlations of outcomes for women within a state, standard errors are clustered by state using a clustered wild bootstrap.

Our outcomes of interest for this procedure are the measures of maternal morbidity and mortality discussed in Section 3, as well as birth rates. We additionally consider exploratory analyses examining post-partum depression. We thus implement the procedure for a measure of all abortion morbidity, morbidity due to haemorrhage early in pregnancy, post-partum depression, and total maternal mortality and maternal mortality due to abortion. In each case in the main outcomes, we focus on rates of morbidity and mortality per the population of fertile aged women. We express our outcomes in this way for two reasons. The first is that it allows us to capture the full effect of the



reform. As we will show that the abortion reform reduces fertility, if we express our outcomes as morbidity or mortality per live birth, this is equivalent to a partial impact, removing any impact of the reform which flows from the ability to avoid undesired, and potentially risky, births.<sup>16</sup> In practice, we are interested in the total impact of the reform, which consists of the reduction in morbidity and mortality due to fewer births, as well as any direct impact the reform may have on the composition of mothers giving birth. Secondly, this allows us to ignore any challenges arising from the endogenous decision of whether or not to engage in legal abortion. If we instead report the impact of the law on rates of morbidity and mortality per live birth, we will be confounding our estimates due to the fact that a non-random group of women choose to proceed with births following the reform, and this group may be selectively more or less healthy than the women who elect to abort. We address changes in the composition of mothers explicitly in Section 5.3 of this paper.

For two-way FE estimates to capture the causal effects of abortion laws, we require a parallel trend assumption to hold, or that outcomes in each of the “Regressive”, “ILE” and untreated states would have evolved similarly in the absence of abortion reforms. We provide a partial test of this, and additionally quantify any dynamic reform effects, by estimating the following panel event-study specifications:

$$Health_{st} = \kappa_0 + \sum_{j=-36}^{36} \delta_{-j} \Delta ILE_{s,t+j} + X'_{st} \Gamma_1 + \phi_s + \mu_t + (\phi_s \times month) + v_{st} \quad (3)$$

$$Health_{st} = \kappa_1 + \sum_{k=-36}^{36} \gamma_{-k} \Delta Regressive_{s,t+k} + X'_{st} \Gamma_2 + \phi_s + \mu_t + (\phi_s \times month) + v_{st}. \quad (4)$$

We normalise both  $\delta$  and  $\gamma$  setting  $\delta_{-1} = 0$  and  $\gamma_{-1} = 0$ . These event-study specifications are increasingly common in panel settings, and here we adopt the notation of Freyaldenhoven et al. (2018). In this specification, we are interested in the leads and the lags of the policy changes, where lags capture any prevailing trends prior to the reform, and leads show the change in health outcomes following the reform’s implementation. In main specifications 3-4, we present the model for morbidity data available from 2001 to 2015, which allows us to consider a large number of monthly lags and leads. We use 36 monthly lags and leads, where the final lag and lead indicates all periods beyond

---

<sup>16</sup>It is also important to note that the ILE reform included the option of accessing free contraceptives after undergoing an abortion procedure, which could also impact birth rates. We discuss this at more length in Section 5.3.

this time. As was the case with two-way FE models, we also examine robustness to using yearly data with all hospitalizations. In this case we are able to use data exclusively from 2004-2015, aggregated to a yearly-level, and as such we estimate 3 leads and 8 lags of the ILE reform, and 5 leads and 7 lags of regressive law changes. We can estimate additional lags for regressive reforms in this case given that there is greater variation in treatment timing.<sup>17</sup> In the case of mortality or fertility where longer periods of data are available, lags and leads are modified in yearly event studies to provide a fully saturated model in each case. In one case where parallel pre-trends appear less convincing, we conduct a newly described estimation and inference procedure known as “Honest DD” to examine possible implications of violations of the parallel trend assumption (Rambachan and Roth, 2019).

It is important to note here that unbiased estimation of reform impacts hinges, pivotally, on the assumption of (conditional) parallel trends, and on a SUTVA assumption implying no spillovers to un-treated states. We address the SUTVA assumption explicitly in Section 5.3 where we consider whether we observe appreciable policy spillovers. As always, the parallel trends assumption is something that cannot be tested formally given the unobserved counter-factual state for treated areas in the post-treatment period. Thus, the appearance of any simultaneous reforms *only* in treatment areas would result in biased estimates of reform impacts. Nevertheless, as discussed above, we aim to reduce the likelihood of such factors by controlling for time-varying factors (including the *Seguro Popular* rollout), and additionally consider a range of placebo tests to determine whether we observe similar patterns in variables which plausibly should *not* respond to abortion reforms.

## 5 Results

### 5.1 Abortion Laws and Maternal Morbidity and Mortality

#### 5.1.1 Estimated Impacts on Maternal Morbidity

Two-way FE estimates (which for simplicity hereafter we will refer to as “DD” estimates given the similarity to double difference models) of the impact of the legal reforms on morbidity are presented in Table 2. All coefficients are cast as the effect of law changes on morbidity per 1,000 women. Columns 1-2 are baseline DD models including only time and state fixed effects, while columns 3-4 add in time-varying controls described previously. We present results for equation 1 in Panel

---

<sup>17</sup>Practical details are discussed in Clarke and Tapia Schythe (2020).

A, and for equation 2 in Panel B. On average, compared with non-reform states, the ILE reform resulted in a reduction in morbidity by approximately 0.06 to 0.08 cases per 1,000 women (per month) when considering all abortion-related morbidity, or be 0.013 to 0.016 cases per 1,000 women when considering the incidence of haemorrhage early in pregnancy. When compared to average rates of morbidities, this is approximately a 20% reduction in abortion related morbidity, and a 35% reduction in rates of haemorrhage. Note that these large effects were notable even in raw trends displayed in Figure 1.

In the case of subsequent restrictive reforms, we do not find evidence to suggest that these reforms shifted morbidity outcomes. For abortion related morbidity and for haemorrhage early in pregnancy, we find no significant impacts across specifications reported in Table 2. When instead of the total number cases we examine the total number of inpatient days (Appendix Table A6), we similarly observe a large reduction following Mexico's ILE reform, and no significant, or consistently signed, impact in the case of regressive reforms.


Figure 2a presents event study evidence for haemorrhage, and Figure 2b presents event study estimates for abortion-related morbidity. In both cases, the left-hand panel shows the lag and lead coefficients for Mexico DF surrounding the ILE reform (equation 3), and the right-hand panel shows the coefficients for regressive states (equation 4). In Figure 2a we observe an immediate sharp decline in rates of haemorrhage in Mexico DF following the adoption of ILE. Additionally, we observe little evidence of prevailing differences in treated and untreated states *before* the reform in all lags. In the case of regressive states (panel B), we observe a similar quite flat profile prior to the reform. Following the reform, we observe no similar reduction in rates of haemorrhage as that observed in DF, with no lead terms being statistically distinguishable from zero at 95% significance levels. These results support claims from the medical literature that haemorrhage is one of the major drivers of maternal morbidity and mortality following unsafe abortions (World Health Organization, 2011), as the appearance of a legal and sterile alternative to clandestine abortion resulted in an immediate sharp reduction in hospitalisations resulting from haemorrhage early in pregnancy. What's more, we observe that these reductions occur very quickly following the moment that abortion was legalised, suggesting an immediate effect of safe legal abortion on women's health outcomes.

Table 2: Difference-in-Differences Estimates of Legal Reforms on Morbidity


	Abortion Related Morbidity			Haemorrhage Early in Pregnancy				
	(1)	(2)	(3)	(4)	(5)	(6)	(7)	(8)
<b>Panel A: ILE versus Non-Reformers</b>								
Post-ILE Reform (DF)	-0.064*** (0.013)	-0.071*** (0.006)	-0.063*** (0.016)	-0.075*** (0.012)	-0.014*** (0.004)	-0.016*** (0.004)	-0.014* (0.007)	-0.013*** (0.007)
Observations	2,496	2,496	2,496	2,496	2,496	2,496	2,496	2,496
Mean of Dependent Variable	0.338	0.338	0.338	0.338	0.044	0.044	0.044	0.044
<b>Panel B: Regressive Reforms versus Non-Reformers</b>								
Post-Regressive Law Change	-0.001 (0.013)	0.000 (0.011)	-0.009 (0.011)	-0.007 (0.011)	-0.005 (0.006)	-0.003 (0.004)	-0.005 (0.006)	-0.002 (0.004)
Observations	5,952	5,952	5,952	5,952	5,952	5,952	5,952	5,952
Mean of Dependent Variable	0.328	0.328	0.328	0.328	0.042	0.042	0.042	0.042
State and Year×Month FEs	Y	Y	Y	Y	Y	Y	Y	Y
Population Weights		Y		Y		Y		Y
Time-Varying Controls			Y	Y			Y	Y

Notes: Each column displays a difference-in-differences regression of the impact of abortion reform on rates of morbidity (inpatient cases). Each morbidity class is measured as cases per 1,000 fertile aged women each month, and average levels in the full set of data are available at the foot of the table. Each regression is estimated using states that adopt reforms (ILE in panel A, regressive reforms in panel B) versus other non-adopting states. All standard errors are clustered at the level of the state. \*p< 0.10; \*\*p< 0.05; \*\*\*p< 0.01.

Figure 2a: Event Studies for Rates of Haemorrhage Early in Pregnancy




(a) Progressive Abortion Reform (ILE)




(b) Regressive Abortion Laws (Legal Tightening)

Figure 2b: Event Studies for Rates of Abortion Morbidity



(c) Progressive Abortion Reform (ILE)



(d) Regressive Abortion Laws (Legal Tightening)

Notes: Event studies document the evolution of rates of haemorrhage early in pregnancy (2a) and abortion related morbidity (2b) per 1,000 women surrounding the passage of abortion reforms. Each point estimate refers to the change in rates between treated and non-treated states, compared to their baseline differential immediately prior to the reform. The left-hand panel shows the difference between Mexico DF and untreated states surrounding the passage of the ILE reform. The right-hand panel shows the difference between regressive policy changers and non-changers around the (time-varying) date that each reform was passed. Regressions are weighted by the population of fertile-aged women, and the full set of time-varying controls are included.

When considering rates of abortion morbidity, event studies document larger prevailing (pre-reform) differences between DF and untreated states, although these are largely driven by a number of monthly fluctuations greater than 2 years prior to the reform.<sup>18</sup> This agrees with simple trends in

<sup>18</sup>In Appendix Figure A5 similar results, with fewer fluctuations in pre-periods, are observed when considering trimesterly rather than monthly cells.

outcomes documented in Appendix Figure 1, which suggest a number of relatively sharper jumps observed in Mexico DF prior to 2005 that were not seen in the rest of the country. However, these *do not* seem to point to radically different pre-reform trends, but rather cyclical outliers. In the case of states which altered legislation in response to ILE, we observe very little evidence of an impact of these reforms on abortion morbidity in 2b. In both the pre- and post-reform period, all estimates are not statistically distinguishable from zero, and are centred around a null impact. In the case of abortion morbidity, it is important to note that the procedure used for abortions realized under the auspices of ILE has changed over time, which may partially explain the delay in observed impacts on morbidity. Initially, the majority of abortions were performed by surgical procedures compared to medical abortions (25%). This gradually changed in subsequent years, with medical abortion procedures reaching 74% in 2011, and the use of dilation and curettage was eliminated entirely (in accordance with WHO recommendations for first-trimester abortions). In addition, the quality of medical abortions performed has also improved, due to the introduction of mifepristone (combined with misoprostol) in 2011 (Becker, 2013).

We note that these reductions in rates of maternal morbidity in Mexico DF surrounding the passage of the ILE reform are not simply capturing a general improvement in health outcomes in the region. In Appendix Figure A6 we estimate identical models for (late-term) obstetric complications, and observe no such improvements in Mexico DF, and evidence to suggest parallel trends throughout the entire period studied between Mexico DF, states implementing regressive reforms, and the rest of the country. Similarly, in Appendix Figures A7 and A8, when considering a much broader class of morbidity outcomes (such as neoplasms, endocrine diseases and so forth), we once again observe no systematic difference in rates of morbidity surrounding abortion reforms, with the exception of a number of monthly peaks/troughs observed in Mexico DF, indicative of no general improvements across the time period studied.

### **5.1.2 Maternal Mortality and “The Tip of the Iceberg”**

Frequently, analyses of the impact of public policies on maternal health focus on maternal mortality, given a lack of access to high-quality morbidity records (such as those available in Mexico). An argument is made that if effects are observed on maternal morbidity, which is the “tip of the

iceberg” (e.g. Firoz et al., 2013), then there will logically be considerable impacts on maternal morbidity (the base of the iceberg). However, the much lower frequency of maternal deaths compared with maternal morbidity makes it much harder to precisely estimate impacts of health reforms on maternal mortality. Here we briefly consider how much precision is lost when considering impacts on maternal mortality, comparing our previously estimated impacts on morbidity with those focusing on maternal deaths. Given the much lower rates of maternal deaths, we focus on yearly models. And as these estimates are indeed considerably more noisy than estimated impacts on morbidity, we relegate results to an Online Appendix, discussing here broad patterns and lessons for estimating reform impacts on maternal health outcomes.

In Appendix Table A7 we present DD estimates following equations 1 (panel A) and 2 (panel B) for both all maternal deaths (columns 1-4), and only maternal deaths originating from abortive causes (columns 5-8). When focusing on the ILE reform, we observe mixed evidence pointing in the direction of negative though imprecisely estimated effects. Both weighted and unweighted estimates suggest a reduction in all maternal deaths following ILE (columns 1 and 2), of approximately 0.5 per 100,000 fertile aged women (versus a mean value of 4 deaths per 100,000 women in Mexico). Note however, that when adding time-varying controls in column 3, these estimates are reduced by about one third, and become statistically insignificant at typical levels. Similarly, in the case of abortion related maternal mortality, we observe significant reductions when using weighted or unweighted simple DD models (with point estimates of around -0.09 per 100,000 fertile aged women), though these become insignificant with the inclusion of time-varying controls. In the case of regressive reforms we find, across the board, relatively less evidence of any impacts of these reforms on maternal mortality. We do consistently observe negative point estimates of a magnitude approaching that observed in Mexico DF following the ILE reform, however these are only (marginally) significant in two models. We note however that, as discussed, standard errors are quite wide, thus precluding us from concluding that these estimates suggest tightly estimated zero-impacts.

In general these results point to the fact that focusing only on mortality when studying reforms which impact maternal health may considerably understate their importance as a determinant of well-being. Both Table A7 (as well as event studies presented in Appendix Figures A9a for all maternal mortality and A9b for mortality due to abortion) suggest noisy results with little power to reject a range of nulls. While event studies suggest that reductions in mortality may have been observed in

Mexico DF, these are certainly not as clearly observed as the morbidity results presented in Section 5.1.1 of this paper.

## 5.2 Benchmarking Impacts on Birth Rates

While our main focus in this paper is on quantifying the health costs of abortion reform, it is illustrative to also estimate impacts on birth rates. This allows us to consider the magnitude of these reforms compared with a range of other contexts which have been well-documented in the economic literature. As summarised in Table A8, across studies on abortion legalization in the US, Nepal, Norway, Romania and Uruguay we observe a drop in birth rates of between 1.2 to 8%. Studies on the impact of regressive abortion law changes (including parental consent laws and restricted funding of abortions) find considerably more heterogeneous results, with results ranging from significant reductions in birth rates (Kane and Staiger, 1996), insignificant impacts (Levine et al., 1996), and increases in rates of birth (Lahey, 2014).

Table 3: Monthly Difference-in-Differences Estimates of Abortion Reforms on Fertility

	Births per 1,000 Women			
	(1)	(2)	(3)	(4)
<b>Panel A: ILE versus Non-Reformers</b>				
Post-ILE Reform (DF)	-0.531*** (0.075)	-0.636*** (0.106)	-0.537*** (0.110)	-0.608*** (0.106)
Observations	2,028	2,028	2,028	2,028
Mean of Dependent Variable	7.302	7.302	7.302	7.302
Mean of Dependent Variable (Mexico DF)	7.433	7.433	7.433	7.433
<b>Panel B: Regressive Reforms versus Non-Reformers</b>				
Post Regressive Law Change	-0.191** (0.089)	-0.269*** (0.097)	-0.202** (0.085)	-0.228** (0.112)
Observations	4,836	4,836	4,836	4,836
Mean of Dependent Variable	7.266	7.266	7.266	7.266
Mean of Dependent Variable (Regressive States)	7.434	7.434	7.434	7.434
State and Year FEs	Y	Y	Y	Y
Population Weights		Y		Y
Time-Varying Controls			Y	Y

Each column displays a difference-in-differences regression of the impact of abortion reform on birth rates. Birth rates are measured as the number of births per 1,000 fertile aged women each month. Time-varying controls are documented in Section B.2. All standard errors are clustered at the level of the state using a wild clustered bootstrap procedure.

\*p< 0.10; \*\*p< 0.05; \*\*\*p< 0.01.



Our results from the Mexican abortion reforms suggest broadly similar impacts on birth rates to those observed in other settings following the elimination of abortion restrictions. In Table 3 we present DD estimates of the impact of reforms on birth rates, weighted and unweighted by the number of fertile-aged women in each state by month cell. In general, across specifications, results are quite stable in suggesting a significant reduction in births in Mexico DF following the ILE reform. Depending on estimation weights, we observe a reduction of between 0.53 and 0.64 births per 1,000 women per month, or a reduction of between 7 and 8.5% in birth rates compared with pre-reform levels in the state. Our preferred estimates are those including population weights with full time-varying controls, which suggest a reduction of 0.64 births per 1,000 women of fertile age in the years following the ILE reform, or a 8.5% reduction in birth rates in Mexico DF. In the case of states passing regressive laws altering their penal codes or state constitutions related to abortion, we find some evidence to suggest smaller reductions in birth rates in these DD specifications. Depending on the model, point estimates vary from -0.19 to -0.27 births per 1,000 women per month, or a 2.5 to 3.6% reduction in birth rates.<sup>19</sup> We return to examine the nature of these legal reforms in more detail in Section 5.3, revisiting the smaller estimated impacts on birth rates.


We provide full event studies corresponding to the passage of progressive and regressive reforms in Figure 3a. Given the considerable seasonality, and even monthly variations, in the quantity of births, we additionally present event studies by trimester in Figure 3b which allow for some smoothing of sharp monthly changes. In the left-hand panel of both monthly and trimesterly event studies, we observe a reduction in birth rates in Mexico DF when compared with all non-reform states, which becomes consistently observed from around 7-10 months (or 2-3 trimesters) post-reform (2008) onwards. This is in line with lags in birth rates expected to be observed approximately 7-9 months following the passage of abortion reforms due to the gestational period and limits on gestational length when undertaking abortion. Estimates in the pre-reform period are not completely flat, if anything suggesting evidence of a slightly upward trend a number of years pre-adoption in Mexico DF. If this upward trend were expected to continue in the post-reform period, this would suggest

---


<sup>19</sup>As we will discuss below, we note that these estimates are significant only in the case of two-way FE models, and not in event study models. In general, in the case of the regressive reforms which are rolled out in a time-varying way, two way FE models potentially mis-estimate the true nature of the ATT, and so we find event study estimates more credible. Nonetheless, we do note that in some literature discussed in Table A8 reductions in fertility are observed following legal restrictions on abortion access, and this could potentially be explained if increasing sanctions on abortion act to discourage marginal births. A theoretical model of such a case is provided in Ananat et al. (2009).

that we actually underestimate the true impact of the ILE policy.

Figure 3a: Monthly Event Studies for Birth Rates




(a) Progressive Abortion Reform (ILE)




(b) Regressive Abortion Laws (Legal Tightening)

Figure 3b: Trimesterly Event Studies for Birth Rates



(c) Progressive Abortion Reform (ILE)



(d) Regressive Abortion Laws (Legal Tightening)

Notes: Event studies document the evolution of birth rates per 1,000 women surrounding the passage of abortion reforms. Each point estimate refers to the change in rates between treated and non-treated states, compared to their baseline differential immediately prior to the reform. Figure 3a presents event studies based on monthly birth rates while Figure 3b is based on trimesterly birth rates. In each case, the left-hand panel shows the difference between Mexico DF and untreated states surrounding the passage of the ILE reform. The right-hand panel shows the difference between regressive policy changers and non-changers around the (time-varying) date that each reform was passed. Regressions are weighted by the population of fertile-aged women, and the full set of time-varying controls are included.

This can be formally examined using “Honest DD” methods proposed by Rambachan and Roth (2019). These methods propose a robust estimation and inference technique assuming that trends in the post-event period do not diverge ‘too much’ from those in the pre-event period. We document how our estimated event study coefficients would vary under a range of assumptions using these

“Honest DD” techniques in Appendix Figures A10, A11 and A12. Given that these require bounding *each* figure in the event study, we conduct these methods with yearly analogues of the monthly event studies (confidence intervals for baseline yearly event studies are presented in blue in these plots). Importantly, as well as allowing for varying violations of parallel trends, they document that in each case, if trends had remained constant in the post-event period, the true estimate would be both more negative and consistently statistically significant (as documented in cases where  $M = 0$  in Appendix Figures). In Appendix C we also document the stability of these estimates when compared to a judiciously chosen synthetic control group. The right-hand panel of Figure 3a documents the same point estimates and standard errors for states altering their constitutions or criminal codes to increase legal sanctions on abortion. In the case of regressive reforms, while we observe a gradual reduction in estimated coefficients following legal changes, these effects are not observed to be statistically distinguishable from zero in event study models. In this case, prevailing trends in the pre-reform period are observed to be somewhat downward sloping, with little evidence of a trend break following the implementation of regressive reforms. These event studies suggest that if anything, DD estimates presented in Table 3a likely over-state the impact of legislative tightenings. It is important to note that, given recent advances in work on interpretation of two-way fixed effect models, in the case of regressive abortion laws which are rolled out in a time-varying ways, these event study models which suggest insignificant effects should be preferred over the models presented in Table 3. This owes to the fact that the weighting in the two-way fixed effect models may result in estimates which are not truly capturing the ATE. We return to this point in the following Sub-section.

## 5.3 Understanding impacts of abortions laws

### 5.3.1 Why do we observe changes in women’s health?

In Section 2.4 we discussed three conceptual channels through which abortion reform could impact maternal health: the “quality of care” channel, the “demand” channel, and the “selection” channel. We consider which of these various channels could explain the observed impacts of abortion reform on women’s health laid out in Section 5.1.

The selection channel suggests that the composition of women giving birth may change owing due to selective interaction with abortion laws. We consider this directly in Appendix Table A9

where we use the same two-way fixed effect set-up to examine how maternal/paternal characteristics vary surrounding both the ILE reform as well as regressive reforms. We observe in general, that parents who give birth following the ILE reform are both older, as well as more educated. We *also* see some change of composition of parents following regressive reforms with parents being more likely to be married, potentially suggesting that there are impacts on the types of women conceiving when sanctions on abortions are raised. While this suggests that a *selection* channel may be occurring, it is an indirect test, not allowing us to sign the direction this channel works in, as it does not directly consider the interaction of these changing characteristics on health outcomes. We test this directly in Appendix Table A10. Here we consider the same two-way fixed effect models examining reform impacts on health outcomes, however now additionally controlling for variables which capture parental selection (namely, all outcomes considered in Appendix Table A9). This allows us to ask whether the estimates of the impact of abortion reforms on health that we observe are completely explained by the changes in composition of parents.

The results of Table A10 suggest a number of important interpretations. Firstly, in the case of abortion related morbidity, much of the observed impacts of the ILE reform *do* appear to flow from changes in the characteristics of parents. And in particular, the fact that our significant estimates are attenuated towards zero conditional on these controls suggests that those giving birth following the reform are selectively healthier (at least considering the likelihood of having a miscarriage or other types of abortion related morbidity). However, in the case of haemorrhage early in pregnancy, we observe no such change in the coefficients. This suggests that the observed impacts are not simply a compositional impact, but rather likely owe to the demand channel or the quality of care channel. In the case of Mexico, it seems likely that the quality of care channel explains the large reduction in haemorrhage. In clandestine settings, where misoprostol is used without adequate access to information, haemorrhage and resulting hospitalisation is likely to occur, something which can be avoided when care is sought through ILE providers who give both treatment as well as information related to post-abortion care. However, without data on the usage of clandestine abortion prior to the ILE reform, it is difficult to assess the degree to which legalizing abortion may have resulted in the avoidance of marginal births. Thus, in principle, while the evidence suggests that the observed impacts of legalized abortion on rates of haemorrhage early in birth do not owe to the selection channel, and it seems likely that the quality of care channel is a main determinant, without data on clandestine

abortion, we cannot formally rule out the demand channel.<sup>20</sup>

### 5.3.2 *De Jure versus De Facto Legal Reforms*

In general we find no impact of regressive law changes on resulting morbidity, and small impacts on birth rates. One potential explanation of this is that the although *de jure* changes were made to state constitutions, the *de facto* implementation of laws and penal codes was unchanged. As we document in Appendix Table A1, in many cases, while constitutions were altered—generally to declare that human life begins at conception—this did not always translate in concrete legal changes in the criminal sanctions imposed on women. This has been similarly noted in legal analyses of the reform (Singh et al., 2012b). And even in cases where criminal sanctions were increased, it may be the case that state-level judiciaries do not alter the likelihood of imposing sanctions on abortion.

We examine whether there is evidence of changes in the likelihood of being sentenced to prison for undertaking an abortion, or in the length of prison sentences received, based on the passage of the abortion laws examined in this paper. These data cover all individuals in the country, and here our outcomes are focused explicitly on whether an individual is sentenced to prison for undertaking an abortion. DD results following specifications 1-2 are displayed in Table 4. We observe, firstly, that there is a sharp reduction in the number of prison sentences for undertaking an abortion in Mexico DF following the reform (in line with the legalisation of abortion), however this was observed alongside an estimated increase in sentence lengths.<sup>21</sup> Importantly, we observe evidence of a dual impact in regressive states. We observe mixed evidence pointing to a slight reduction in the number of prison sentences handed down, falling by 1.36 cases in weighted regressions (compared with a mean number of sentences per state and year of 1.806). In the case of the length of sentences, we observe a considerable increase, of between 4.1 and 5.2 years, depending on the specification estimated. In the case of weighted estimates, we observe an average increase of 5.2 years, which is considerable, even at the lower end of the 95% confidence interval, when compared with the mean sentence length

---

<sup>20</sup>Given the expansion of the medical abortion regime (i.e. the use of misoprostol and mifepristone) in Mexico City (Becker, 2013), complications due to the abortion procedures themselves following the ILE reform appear low (Grimes et al., 2006a). It is therefore unlikely that channel 2b, from Section 2.4, explains large movements in maternal health.

<sup>21</sup>Note that in Mexico DF, while abortion was legalised by the ILE reform, this was only the case for abortions realised up to 12 weeks of gestation. Thus, in theory, custodial sentences can still be handed down for abortion when not meeting this condition. In practice, a non-zero number of sentences was only observed in Mexico DF in 2011 (refer to Appendix Figure A13 for trends over time).

Table 4: Difference-in-Differences Estimates of Abortion Reforms on Judicial Outcomes

	Number of Prison Sentences		Length of Prison Sentences	
	(1)	(2)	(3)	(4)
<b>Panel A: ILE versus Non-Reformers</b>				
Post-ILE Reform (DF)	-4.018*** (0.294)	-4.050*** (0.340)	2.251** (1.032)	2.150** (1.063)
Observations	117	117	56	56
Mean of Dependent Variable	1.427	1.427	3.606	3.606
<b>Panel B: Regressive Reforms versus Non-Reformers</b>				
Post-Regressive Law Change	-0.648 (0.464)	-1.362** (0.607)	4.144* (2.361)	5.220** (2.439)
Observations	279	279	171	171
Mean of Dependent Variable	1.806	1.806	3.749	3.749
State and Year FEs	Y	Y	Y	Y
Population Weights		Y		Y

Difference-in-difference models illustrate how abortion reforms correlate with prison sentences handed down by the judiciary, and the length of these prison sentences in years. Total number of sentences and the average length of prison sentences are generated from administrative records captured in Mexico's Judicial Statistics on Penal Matters. This is the universe of judiciary decisions in the country based on the first legal judgment, and so does not include any subsequent appeals. Analysis of the length of prison sentences presented in columns 3 and 4 is conditional on any prison sentences being handed down in each state and year. Prison sentence lengths are calculated from a categorical variable capturing bins of between 6 months and two years, and in each case we record the total years (or fractions of years) based on the midpoint of each bin. Bins are consistently used in the period displayed here. All standard errors are clustered at the level of the state and calculated using a wild bootstrap procedure. Identical models using population-standardised values for dependent variables are presented in Appendix Table A11. \* $p < 0.10$ ; \*\* $p < 0.05$ ; \*\*\* $p < 0.01$ .

of 3.7 years. Thus, these results suggest that while the changes in law did not necessarily always prescribe a change in prison sentences, there is a detectable increase in the length of prison sentences observed in administrative data, conditional on being sentenced to prison. This increase in average sentence length is observed to hold in event study analysis, with significant impacts observed from 1 year post-reform onwards (Appendix Figure A14).

### 5.3.3 Considering the Universe of Health Records

**Considering the Universe of the Public Health System** In principal models, we have used month  $\times$  year  $\times$  state measures of outcomes, given that as documented in Table A2, a wave of legal restric-

tions were put in place in varying months of 2008 and 2009. However, as discussed in the Data Section of this document, only hospitals administered by Mexico’s Secretariat of Health (available to all individuals) report month of hospitalisation in administrative data. In order to determine whether a similar pattern is observed across the entire public health system, we additionally conduct analysis pooling by hospitalisations in hospitals administered by the Secretariat of Health, as well as those administered by Social Security Providers. In this case, given the lack of monthly records in Social Security Hospitals, we can only estimate models by year. In Figure 4 we present yearly event studies for the main morbidity outcomes considered, extending to this full sample.

In all cases, we observe broadly similar results in each case. In the case of haemorrhage morbidity, we observe a sharp fall of around 1 per 1,000 cases per year immediately following the reform in Mexico DF (panel a), with no similar results observed states undertaking reform tightenings (panel b). Similarly, reductions in abortion related morbidity are observed in Mexico DF, appearing gradually over time as observed in monthly event studies, with no similar changes observed in regressive states (panels (c) and (d)).


**Leakage to the Private Health System** One potential alternative explanation of the observed morbidity results in all public hospitals is that rather than being driven entirely by the abortion reform, they may reflect changes of usage of the health system, with a larger number of women opting to use the private health care system. This explanation cannot explain the impact on fertility and maternal mortality, as these outcomes are based on the complete records of births and deaths in the country. However, it could partially explain the impacts observed on morbidity, as our administrative data records inpatient stays in the public health system (refer to Section 3 for additional discussion).

While we can’t consistently merge public and private health data at the most disaggregated level of morbidity causes, we *are* able to consider all causes of abortion morbidity in the private health care system.<sup>22</sup> In Appendix Figure A15 we plot rates of abortion related morbidity in the universe of private hospitals (left-hand panel) and the universe of public hospitals (right-hand panel). These descriptive plots suggest that if anything, results in the private system will only strengthen our esti-


---

<sup>22</sup>Note that as documented in Appendix Table A3, this mapping captures all ICD-10 codes O00-O08, while typically abortion morbidity is calculated from codes O02-O08. In Figure A15 we plot comparisons using precisely the same aggregated codes in public and private hospitals.


Figure 4: Yearly Event Studies Based on all Public Data




(a) Haemorrhage (Progressive)



(b) Haemorrhage (Regressive)



(c) Abortion Related (Progressive)



(d) Abortion Related (Regressive)

Notes: Event studies replicate those from Figures 2a and 2b, however using yearly administrative records based on Secretariat of Health hospitals *and* Social Security Hospitals. All details follow those indicated in notes to Figures 2a and 2b however we now work with the universe of hospital visits in all public hospitals. DD models are provided in Appendix Table A12 (and Appendix Table A13 for birth rates).

mates, as we observe a sharper reduction in abortion related morbidity in private hospitals than we observe in public hospitals. In the case of morbidity due to haemorrhage early in pregnancy, we are unable to observe this as a sole cause in the private health records, but we are able to observe the class in which this cause falls and again we observe a considerably sharper reduction in morbidity following the reform in the private health system than we observe in the public health system (Appendix Figure A16).



### 5.3.4 Reform Spillovers and Heterogeneity

**Reform Spillovers** As outlined in Section 2.2, the ILE reform was not strictly limited to residents of Mexico DF. Recent evidence from the United States documents a willingness to travel over a significant distance to access abortion providers (Lindo et al., 2019; Venator and Fletcher, 2019). In Appendix Table A14 we provide summary figures of the state of precedence of users of abortion services in Mexico DF based on administrative data for 2007-2015. While the majority of users (72.5%) are women from Mexico DF, women residing all throughout Mexico have access to ILE. The largest non-DF population comes from nearby Mexico State (24.2%). In general, users of the ILE reform are clustered in states geographically close to Mexico DF. A descriptive plot is presented in Appendix Figure A17. Residents in Mexico DF have by far the highest rate of abortion, at 5.8 abortions per 1,000 women aged 15-49, followed by Mexico State (at 1 per 1,000), and then two nearby states (Hidalgo and Morelos) with rates of 0.1 per 1,000. Remaining states have rates which are an additional order of magnitude lower than this.

Despite some evidence of very localised geographic spillovers, we do not observe clear evidence of changes in birth or maternal health outcomes in nearby states. In Table A15 we estimate DD models augmenting specification 1 with a post ILE $\times$ spillover state indicator, where spillover states refer Mexico State, Morelos and Hidalgo (the three states with most considerable abortion usage per population). In no case do we observe statistically significant reductions in fertility, morbidity or mortality, if anything observing weakly positive impacts. Additional discussion is provided in Appendix C.

An alternative model which captures both the impacts of the reform in Mexico DF as well as any reform spillovers to the rest of the country, replaces the ILE variable in equation 1 with the intensity of treatment in each state. This intensity measure is captured as the rate of abortion per 1,000 women (documented in Appendix Table A14) in the post-reform period in each state. This information is reported in official ILE reports, however is only available at the level of the year, and as such in this case we estimate yearly models as in Section 5.3.3. If outcomes per 1,000 women are regressed on abortion usage per 1,000 women, this provides a back-of-the-envelope calculation of the elasticity of outcomes with respect to the availability of a legal abortion. For example, if each additional legal abortion results in 1 fewer births, we will estimate a coefficient of  $-1$  in this model, suggesting full

pass-through of legalised abortion to birth rates. We note however in our case that the figures on abortions refer *only* to those abortions provided by official ILE providers. To the degree that private providers additionally provide access to legal abortion, these estimates should be seen as an upper bound estimate on the magnitude of pass through. We estimate models of this type in Appendix Table A16. In general we observe that, using the full data on abortions across Mexico resulting from the ILE reform, impacts per abortion are considerable, suggesting approximately 0.9 fewer birth per every abortion provided, 0.16 fewer cases of abortion related morbidity, and 0.14 fewer cases of morbidity due to haemorrhage. These values must be viewed in line with the caveat above, that any abortions accessed from private providers will reduce the actual magnitude of pass-through (likely considerably given the relevance of the private sector), and as such, these are upper bound estimates.

**Individual Heterogeneity in Reform Impacts** Effects discussed up to this point have focused on average outcomes over all women aged 15–49. However, given evidence documenting heterogeneous impacts of abortion reform in other contexts (Ananat et al., 2007), and heterogeneity in access to abortion in Mexico (Friedman et al., 2019), we consider heterogeneity in a number of contexts here. In Tables A17 and A18 we examine the impacts of the reform on abortion related morbidity and on haemorrhage early in pregnancy by quinquennial age groups and by individual’s insurance coverage. While we would like to consider impacts by individual income level, this latter category of insurance coverage is one of the closest proxies of income which we can observe in hospitalization microdata. In the case of both outcomes we observe three broad stylised facts. The first, health impacts are observed across the age distribution; the second, that these impacts are largest in size among younger women, peaking in the age group of 20–24 years; and the third that results appear to be driven by individuals who *do* have insurance coverage. This final fact suggests that impacts may be higher among higher income women, however it is important to note that this classification by insurance status is very crude, as not having formal insurance likely signifies a considerable disconnection from the public health system.

Finally, in Appendix Figure A18 we document impacts more finely by age, specifically focusing on teen-aged and younger women. Given that minors require parental consent, we may expect that impacts on these women are reduced, in line with reduced rates of usage. In Appendix Figure A18 we observe single-coefficient DD estimates for each age, where we estimate models 1 (presented


as hollow circles) and 2 (presented as black squares) separately in each yearly age group. Here, while we do observe a general increase in the magnitude of coefficients while increasing from 15 to 18 years and above, we do observe that even among younger adolescents (for example 16 year-olds), significant impacts of ILE are observed on both rates of haemorrhage, and abortion related morbidity.

**Temporal Heterogeneity in Reform Impacts** An additional concern related to the timing of adoption of the regressive reforms is that single-coefficient DD estimates (equation 1) may be capturing heterogeneity and variation in law implementation, rather than the ATT itself (Goodman-Bacon, 2018). In particular, we may be concerned that small effects are being driven by the incorrect use of already treated units as controls in future periods. This will bias towards zero any effects of the laws if these impacts are growing over time. We thus follow Goodman-Bacon (2018) in explicitly decomposing the single-coefficient DD model into its component parts of a pure “treated versus never-treated” effect, and effects owing simply to the variation in timing of the passage of laws. We note that this decomposition is only of concern in the case of regressive laws given that ILE was adopted in a single moment of time (and a single state). This decomposition is displayed graphically in Figure 5 following Goodman-Bacon (2018); Goodman-Bacon et al. (2019). We plot the full set of “ $2 \times 2$ ” DD estimates of the impact of regressive reforms on morbidity outcomes,<sup>23</sup> where these models come from all variation in treatment timing and all possible control groups. We observe that estimates are largely clustered around zero, particularly for the Treated versus Never-Treated comparisons of interest. We present the global decomposition in Appendix Table A19, and observe—reassuringly—that in each case the majority of the weight in the single coefficient DD estimate comes from the Treated versus Never-Treated comparison (around 92%), and in general, estimates even within the timing-only groups are similarly small when compared to the Treated versus Never-Treated effect.

---

<sup>23</sup> Estimates for birth rates are provided in Appendix Figure A19.

Figure 5: Goodman-Bacon (2018) Decomposition Based on  $2 \times 2$  Difference-in-Difference Models



Notes: Figures document the Goodman-Bacon decomposition into a series of  $2 \times 2$  difference-in-differences models depending on the type of comparison unit. Here the “treatment” refers to the passage of a regressive abortion law, and the outcomes refer to rates of morbidity per 1,000 women. The passage of the ILE reform occurred at a single moment in time, and as such, decompositions need not be performed. The global decomposition is given in Table A19.

### 5.3.5 Broader Impacts of the ILE Reform


The ILE reform was a significant change, noteworthy even within the entire continent of Latin America (Sánchez Fuentes et al., 2008). Evidence from literature in economics such as Chiappori and Orefice (2008); Orefice (2007) suggests that abortion reform can have significant impacts beyond health and fertility outcomes, extending to broader spheres within households and individual well-being. Here we briefly discuss a number of possible additional impacts of abortion reform on women’s mental health and family outcomes. Given challenges in measurement and reporting,<sup>24</sup> this work should be considered as exploratory, and subject to future research.

To examine the impact on maternal mental health, we consider the predominant mental disorders in this domain—namely post-partum depression and other mental and behavioral disorders associ-

<sup>24</sup>When screened for, maternal depression (and other mental health issues related to childbirth) is a common condition both antenatal and post-partum, with estimates suggesting a prevalence between 13-18% in Mexico (Albuja et al., 2017). However, as in most countries, the prevalence of maternal depression is difficult to establish as maternal depression is often under-diagnosed and under-reported (on the end of both health care professionals as well as patients) (Anokye et al., 2018). Our ability to capture post-partum depression (and similar conditions) is limited given that we don’t have access to outpatient or pharmaceutical data. Being hospitalised for post-partum depression is rare and hospitalisations are likely to capture the most severe cases of mental illness while the majority of women are most likely treated elsewhere if diagnosed.

ated with the puerperium (Organization et al., 2015). We use the same administrative health records as used previously, recording all hospitalisations classified as owing to mental and behavioral disorders associated with the puerperium including post-partum depression (ICD-10 code F53).<sup>25</sup>


Figure 6a: Monthly Event Studies Examining Post-Partum Depression



(a) Progressive Abortion Reform (ILE)

(b) Regressive Abortion Laws (Legal Tightening)

Figure 6b: Yearly Event Studies Examining Post-Partum Depression



(c) Progressive Abortion Reform (ILE)

(d) Regressive Abortion Laws (Legal Tightening)

Notes: Event studies document the evolution of rates of post-partum Depression per 1,000 women surrounding the passage of abortion reforms. Each point estimate refers to the change in rates between treated and non-treated states, compared to their baseline differential immediately prior to the reform. Figure 6a presents event studies based on monthly rates of post-partum Depression (using data from Secretary of Health hospitals only), while Figure 6b presents event studies based on yearly rates of post-partum Depression using data from all public hospitals (both Secretary of Health and Social Security providers). In each case, the left-hand panel shows the difference between Mexico DF and untreated states surrounding the passage of the ILE reform. The right-hand panel shows the difference between regressive policy changers and non-changers around the (time-varying) date that each reform was passed. Regressions are weighted by the population of fertile-aged women, and the full set of time-varying controls are included.

In Figure 6a and 6b we estimate impacts of abortion reform following equations 1-2 on rates of

<sup>25</sup> Another relevant mental health outcome is suicide attempt. However, this is an extreme measure of poor mental health. Moreover, in the ICD system suicide attempts are recorded separately as external causes, which are not consistently available in all micro-data over time.

post-partum depression. While Figure 6a suggests no consistently observable impacts on rates of post-partum depression around the passage of abortion reforms (for both progressive and regressive reforms), there is some evidence suggesting a reduction in rates of depression in Mexico DF compared with other non-treated states when examining yearly estimates in Figure 6b. No such clear evidence is observed in the case of states undertaking legal tightenings.

Finally, while in this paper we are focusing principally on the health impacts of the adoption of abortion laws, there are potentially much broader impacts of such a policy. For example, Lauletta (2019) provides evidence of potential reductions in domestic violence flowing from these reforms, in line with economic theory suggesting improvements in women's bargaining power flowing from birth control reforms.

### 5.3.6 Mechanisms: Availability, education, or behavior

Along with the law change legalising abortions, the ILE reform included additional components relating to sexual education and disbursement of additional contraceptives in clinics (refer to Section 2.2). To examine the channels through which the reform affected birth rates: whether it be only access, or a combination of access with behavioral change, we use the MxFLS data which follows women over time, and has survey rounds both before and after the abortion reforms of interest. To examine the potential effect of the other aspects of the reform (sexual education and contraceptives), we estimate a version of equation 1, however at the level of the individual, which allows for individual-specific fixed-effects given the panel nature of the MxFLS data.

We examine the effect of abortion reform on all available measures of contraceptive use (using any contraceptive or using modern contraceptives), the number of reported sexual partners and whether the respondent reports having knowledge of modern contraceptive methods. These results are presented in Appendix Table A20. In general, we find very little evidence to suggest that the results of the abortion reform flow from an increase in *other* contraceptive knowledge in reform areas, or change in risky sexual behavior as a result of the reform. We find quite close to zero effects for change in contraceptive use and knowledge, and an insignificant reduction in the number of sexual partners reported. In all cases, these results are insignificant at the 10% level<sup>26</sup> suggesting that the

---

<sup>26</sup>Summary statistics are provided in Appendix Table A21, and similar results are observed using a repeated cross

ILE reform's effect is largely due to the sharp increase in utilization of abortion services, rather than alternative contraception or information channels. Similarly, we do not find that regressive changes in abortion laws cause women to seek additional information or be more likely to use contraceptives, or change sexual behavior as proxied by the number of sexual partners compared to areas which were not subject to a regressive reform. We do note in the case of the number of sexual partners that while we cannot reject that the impact is significant even at a 10% level, we cannot rule out economically meaningful effects given the reasonably inexact point estimate.

## 6 Conclusion

In this paper we examine the impact of abortion law on women's health. We consider a context in which considerable heterogeneity in legislative reform is observed. In Mexico in the late 2000s both a substantial loosening, and a series of tightenings of abortion policies were undertaken at the sub-national level. Using comprehensive vital statistics data on maternal health outcomes, we observe that safe legal abortion available in the first trimester of pregnancy in Mexico DF resulted in a sharp drop in maternal morbidity due to haemorrhage, and a slower decline in morbidity due to abortion, perhaps in line with the gradual adoption of recommended abortion techniques by public health clinics. These declines were of substantial importance, suggesting 8,600 fewer inpatient visits in the post-abortion years in Mexico DF. In general, we observe quite weak effects of the tightening of *de facto* sanctions on abortion, even though, as we show, these sanctions did lead to changes in the length of sentences handed down to women.

We document that the impact of Mexico DF's ILE reform on birth rates, at around a 8% reduction, is in line with impacts estimated in other settings, for example the US in the 1970s. We observe generally weaker effects of regressive reforms on birth rates, though note that in the case of Mexico, these state-level reforms may have *reduced* fertility by around 1-2%. Importantly, when examining the impacts of abortion reforms on rates of maternal death, our estimates are considerably noisier than those for maternal morbidity. This is of importance given that a range of papers examining the impact of abortion on women's health limit analyses to maternal death, given a paucity of high-quality health records. Our results suggest that this focus on "the tip of the iceberg" may lead to less convincing results than when focusing on maternal morbidity. While focusing on surviving

---

section of women (Appendix Table A22).

childbirth should be an absolute minimum when designing public policies to protect maternal and women's health, maternal morbidity is of considerable importance when quantifying life-time well-being, and avoiding a considerable health burden leading to chronic conditions.

The results of this paper are becoming relevant once again as a number of countries revisit abortion legislation and attempt to make considerable changes in constitutions and penal codes. Among others, legislative reforms have been undertaken or attempted in Ireland, Argentina, Australia, Chile and New Zealand in 2017-2020 focusing on legalising abortion in certain circumstance, and increasing restrictions have been enacted or proposed in Poland and a number of US states. This paper documents that these policies are likely to have a considerable impact on women's health and well-being.

## References

- Abadie, A., A. Diamond, and J. Hainmueller (2010): "Synthetic Control Methods for Comparative Case Studies: Estimating the Effect of California's Tobacco Control Program," *Journal of the American Statistical Association*, 105, 493–505.
- Albuja, A. F., M. A. Lara, L. Navarrete, and L. Nieto (2017): "Social support and postpartum depression revisited: the traditional female role as moderator among Mexican women," *Sex roles*, 77, 209–220.
- Ananat, E. O., J. Gruber, and P. Levine (2007): "Abortion Legalization and Life-Cycle Fertility," *Journal of Human Resources*, XLII, 375–397.
- Ananat, E. O., J. Gruber, P. B. Levine, and D. Staiger (2009): "Abortion and Selection," *Review of Economics and Statistics*, 91, 124–136.
- Ananat, E. O. and D. M. Hungerman (2012): "The power of the pill for the next generation: oral contraception's effects on fertility, abortion, and maternal and child characteristics," *Review of Economics and Statistics*, 94, 37–51.
- Angrist, J. D. and W. N. Evans (1996): "Schooling and Labor Market Consequences of the 1970 State Abortion Reforms," NBER Working Papers 5406, National Bureau of Economic Research, Inc.
- Anokye, R., E. Acheampong, A. Budu-Ainooson, E. I. Obeng, and A. G. Akwasi (2018): "Prevalence of postpartum depression and interventions utilized for its management," *Annals of general psychiatry*, 17, 18.
- Antón, J.-I., Z. Ferre, and P. Triunfo (2018): "The impact of the legalisation of abortion on birth outcomes in Uruguay," *Health Economics*, 27, 1103–1119.
- Bailey, M. J. (2009): "Erratum and Addendum to "More Power to the Pill: The Impact of Contraceptive Freedom on Women's Life Cycle Labor Supply"," Posted at *The Quarterly Journal of Economics*.
- Bailey, M. J. and J. M. Lindo (2017): "Access and Use of Contraception and Its Effects on Women's



- Outcomes in the U.S,” NBER Working Papers 23465, National Bureau of Economic Research, Inc.
- Becker, D. (2013): “Decriminalization of Abortion in Mexico City: The Effects on Women’s Reproductive Rights,” *American Journal of Public Health*, 103, 590–593.
- Becker, D., C. D. Olavarrieta, S. G. Garcia, and C. C. Harper (2013): “Women’s reports on postabortion family-planning services provided by the public-sector legal abortion program in Mexico City,” *International Journal of Gynecology & Obstetrics*, 121, 149–153.
- Benson, J., K. Andersen, and G. Samandari (2011): “Reductions in abortion-related mortality following policy reform: evidence from Romania, South Africa and Bangladesh,” *Reproductive health*, 8, 39.
- Bitler, M. and M. Zavodny (2001): “The effect of abortion restrictions on the timing of abortions,” *Journal of Health Economics*, 20, 1011–1032.
- Blanco-Mancilla, G. (2011): “Implementation of health policies in Mexico City: what factors contribute to more effective service delivery?” Ph.D. thesis, The London School of Economics and Political Science (LSE).
- Bongaarts, J. (2016): “WHO, UNICEF, UNFPA, World Bank Group, and United Nations Population Division Trends in Maternal Mortality: 1990 to 2015 Geneva: World Health Organization, 2015.” *Population and Development Review*, 42, 726–726.
- Chiappori, P.-A. and S. Oreffice (2008): “Birth Control and Female Empowerment: An Equilibrium Analysis,” *Journal of Political Economy*, 116, 113–140.
- Chowdhury, M. E., R. Botlero, M. Koblinsky, S. K. Saha, G. Dieltiens, and C. Ronsmans (2007): “Determinants of reduction in maternal mortality in Matlab, Bangladesh: a 30-year cohort study,” *The Lancet*, 370, 1320–1328.
- Ciudad de México (2007): “Gaceta Oficial del Distrito Federal,” Gaceta Oficial, No 70, Órgano del Gobierno del Distrito Federal, 26 April, 2007, accessed at [http://web.archive.org/web/20150924024041/http://www.inmujeres.df.gob.mx/wb/inmujeres/decreto\\_por\\_el\\_que\\_se\\_reforma\\_el\\_codigo\\_penal\\_para](http://web.archive.org/web/20150924024041/http://www.inmujeres.df.gob.mx/wb/inmujeres/decreto_por_el_que_se_reforma_el_codigo_penal_para) (consulted April 26, 2016).
- Clarke, D. and H. Mühlrad (2018): “Abortion Laws and Women’s Health,” IZA Discussion Papers 11890, Institute of Labor Economics (IZA).
- Clarke, D. and K. Tapia Schythe (2020): “Implementing the Panel Event Study,” IZA Discussion Papers 13524, Institute of Labor Economics (IZA).
- Consejo Nacional de Población (2012): “Proyecciones de la población de México 2010–2050,” Tech. Rep. Documento Metodológico, CONAPO.
- Contreras, X., M. G. van Dijk, T. Sanchez, and P. S. Smith (2011): “Experiences and Opinions of Health-Care Professionals Regarding Legal Abortion in Mexico City: A Qualitative Study,” *Studies in Family Planning*, 42, 183–190.
- Cook, P. J., A. M. Parnell, M. J. Moore, and D. Pagnini (1999): “The effects of short-term variation in abortion funding on pregnancy outcomes,” *Journal of Health Economics*, 18, 241 – 257.
- Darney, B. G., B. Saavedra-Avendano, and R. Lozano (2017): “Maintaining rigor in research: flaws in a recent study and a reanalysis of the relationship between state abortion laws and maternal mortality in Mexico,” *Contraception*, 95, 105–111.
- Firoz, T., D. Chou, P. von Dadelszen, P. Agrawal, R. Vanderkruik, O. Tunçalp, N. v. Laura A Magee,

- and L. Say (2013): “Measuring maternal health: focus on maternal morbidity,” *Bulletin of the World Health Organization*, 91, 794–796.
- Fischer, S., H. Royer, and C. White (2018): “The impacts of reduced access to abortion and family planning services on abortions, births, and contraceptive purchases,” *Journal of Public Economics*, 167, 43–68.
- Fraser, B. (2015): “Tide begins to turn on abortion access in South America,” *The Lancet*, 383, 2113 – 2114.
- Freyaldenhoven, S., C. Hansen, and J. M. Shapiro (2018): “Pre-event Trends in the Panel Event-study Design,” Working Paper 24565, National Bureau of Economic Research.
- Friedman, J., B. Saavedra-Avendaño, R. Schiavon, L. Alexander, P. Sanhueza, R. Rios-Polanco, L. Garcia-Martinez, and B. G. Darney (2019): “Quantifying disparities in access to public-sector abortion based on legislative differences within the Mexico City Metropolitan Area,” *Contraception*, 99, 160–164.
- Gamboa Montejano, C. and S. Valdés Robledo (2014): “Regulación del Aborto en México: Derecho Comparado de los 31 estados y del Distrito Federal, así como de diversos países en el mundo y estadísticas del INEGI en el tema,” SAPI-ISS 33-14, Subdirección de Análisis de Política Interior.
- Gerdts, C., D. Vohra, and J. Ahern (2013): “Measuring Unsafe Abortion-Related Mortality: A Systematic Review of the Existing Methods,” *PLoS ONE*, 8, e53346.
- Gobierno de Nayarit (2009): “Exposición de Motivos; Reforma y Adición al Artículo 7, Fracción XI, de la Constitución Política del Estado Libre y Soberano de Nayarit y cómputo y Declaratoria de Aprobación de la Reforma,” Periódico Oficial 079, Organo del Gobierno de Nayarit.
- Goldstein, P. and G. Stewart (1972): “Trends in therapeutic abortion in San Francisco,” *American journal of public health*, 62, 695–699.
- Goodman-Bacon, A. (2018): “Difference-in-Differences with Variation in Treatment Timing,” Working Paper 25018, National Bureau of Economic Research.
- Goodman-Bacon, A., T. Goldring, and A. Nichols (2019): “BACONDECOMP: Stata module to perform a Bacon decomposition of difference-in-differences estimation,” .
- Grimes, D. A. (2005): “Risks of mifepristone abortion in context,” *Contraception*, 71, 161.
- Grimes, D. A., J. Benson, S. Singh, M. Romero, B. Ganatra, F. E. Okonofua, and I. H. Shah (2006a): “Unsafe abortion: the preventable pandemic,” *The lancet*, 368, 1908–1919.
- (2006b): “Unsafe Abortion: The Preventable Pandemic,” *The Lancet*, 368, 1908–1919.
- Gruber, J., P. Levine, and D. Staiger (1999): “Abortion Legalization and Child Living Circumstances: Who is the “Marginal Child?”” *The Quarterly Journal of Economics*, 114, 263–291.
- Guldi, M. (2008): “Fertility Effects of Abortion and Birth Control Pill Access for Minors,” *Demography*, 45, 817–827.
- Gutierrez-Vazquez, E. Y. and E. A. Parrado (2015): “Abortion Legalization and Fertility Rates in Mexico,” Unpublished manuscript, University of Pennsylvania, available at <http://paa2015.princeton.edu/uploads/151146>.
- Guttmacher Institute (2012): “Facts on induced abortion worldwide,” .
- Haddad, L. B. and N. M. Nour (2009): “Unsafe Abortion: Unnecessary Maternal Mortality,” *Obstetrics & Gynecology*, 2, 122–126.
- Henderson, J. T., M. Puri, M. Blum, C. C. Harper, A. Rana, G. Gurung, N. Pradhan, K. Regmi, K. Malla, S. Sharma, et al. (2013): “Effects of abortion legalization in Nepal, 2001–2010,” *PloS one*, 8, e64775.

- Instituto Nacional de Estadística y Geografía (2012): “Derecho a la identidad. La cobertura del registro de nacimiento en México en 1999 y 2009,” Tech. rep., INEGI and UNICEF.
- Jewkes, R., H. Brown, K. Dickson-Tetteh, J. Levin, H. Rees, et al. (2002): “Prevalence of morbidity associated with abortion before and after legalisation in South Africa,” *British Medical Journal*, 324, 1252–1253.
- Johnson, T. (2013): “Guaranteed Access to Safe and Legal Abortions: The True Revolution of Mexico City’s Legal Reforms Regarding Abortion,” *Columbia Human Rights Law Review*, 44, 437–476.
- Joyce, T. and R. Kaestner (1996): “The effect of expansions in medicaid income eligibility on abortion,” *Demography*, 33, 181–192.
- Joyce, T., R. Kaestner, and S. Colman (2006): “Changes in Abortions and Births Following Texas’s Parental Notification Law,” *New England Journal of Medicine*, 354, 1031–1038.
- Joyce, T., R.-D. Tan, and Y. Zhang (2013): “Abortion Before and After Roe,” *Journal of Health Economics*, 32, 804–815.
- Juarez, F., S. Singh, S. G. Garcia, and C. D. Olavarrieta (2008): “Estimates of induced abortion in Mexico: what’s changed between 1990 and 2006?” *International Family Planning Perspectives*, 158–168.
- Kahan, R. S., L. D. Baker, and M. G. Freeman (1975): “The effect of legalized abortion on morbidity resulting from criminal abortion,” *American Journal of Obstetrics & Gynecology*, 121, 114–116.
- Kane, T. J. and D. Staiger (1996): “Teen Motherhood and Abortion Access,” *Quarterly Journal of Economics*, 111, 467–506.
- Koch, E., M. Chireau, F. Pliego, J. Stanford, S. Haddad, B. Calhoun, P. Aracena, M. Bravo, S. Gatica, and J. Thorp (2015): “Abortion legislation, maternal healthcare, fertility, female literacy, sanitation, violence against women and maternal deaths: a natural experiment in 32 Mexican states,” *BMJ open*, 5, e006013.
- Kulczycki, A. (2011): “Abortion in Latin America: Changes in Practice, Growing Conflict, and Recent Policy Developments,” *Studies in Family Planning*, 42, 199–220.
- Lahey, J. N. (2014): “The Effect of Anti-Abortion Legislation on Nineteenth Century Fertility,” *Demography*, 51, 939–948.
- Lauletta, M. (2019): “Abortion decriminalization and domestic violence: evidence from Mexico,” Tesis de Maestría en Economía 10908-17085, Universidad de San Andrés. Departamento de Economía.
- Levine, P. B., D. Staiger, T. J. Kane, and D. J. Zimmerman (1999): “Roe v Wade and American fertility,” *American Journal of Public Health*, 89, 199–203.
- Levine, P. B., A. B. Trainor, and D. J. Zimmerman (1996): “The effect of Medicaid abortion funding restrictions on abortions, pregnancies and births,” *Journal of Health Economics*, 15, 555–578.
- Lindo, J., C. Myers, A. Schlosser, and S. Cunningham (2019): “How Far Is Too Far?: New Evidence on Abortion Clinic Closures, Access, and Abortions,” *Journal of Human Resources*.
- Loudon, I. (1992): *Death in Childbirth. An international study of maternal care and maternal mortality 1800-1950*. 1992, Oxford: Clarendon Press Oxford.
- Madrazo, A. (2009): “The evolution of Mexico City’s abortion laws: From public morality to women’s autonomy,” *International Journal of Gynecology & Obstetrics*, 106, 266 – 269.
- Mbele, A., L. Snyman, and R. C. Pattinson (2006): “Impact of the Choice on Termination of Preg-

- nancy Act on maternal morbidity and mortality in the west of Pretoria,” *South African Medical Journal*, 96, 1196–1198.
- Mitrut, A. and F.-C. Wolff (2011): “The Impact of Legalized Abortion on Child Health Outcomes and Abandonment. Evidence from Romania,” *Journal of Health Economics*, 30, 1219–1231.
- Mølland, E. (2016): “Benefits from delay? The effect of abortion availability on young women and their children,” *Labour Economics*, 43, 6–28.
- Mondragón y Kalb, M., A. Ahued Ortega, J. Morales Velazquez, C. Diaz Olavarrieta, J. Valencia Rodriguez, D. Becker, and S. Garcia (2011): “Patient characteristics and service trends following abortion legalization in Mexico City, 2007-10,” *American journal of public health*, 3, 159–66.
- Myers, C. K. (2017): “The Power of Abortion Policy: Reexamining the Effects of Young Women’s Access to Reproductive Control,” *Journal of Political Economy*, 125, 2178–2224.
- Nunes, F. E. and Y. M. Delph (1997): “Making abortion law reform work: steps and slips in Guyana,” *Reproductive Health Matters*, 5, 66–76.
- Oreffice, S. (2007): “Did the legalization of abortion increase women’s household bargaining power? Evidence from labor supply,” *Review of Economics of the Household*, 5, 181–207.
- Organization, W. H. et al. (2015): “Thinking healthy: a manual for psychosocial management of perinatal depression, WHO generic field-trial version 1.0, 2015,” Tech. rep., World Health Organization.
- Pazol, K., A. A. Creanga, K. D. Burley, and D. J. Jamieson (2014): “Abortion Surveillance – United States, 2011,” Tech. rep., Division of Reproductive Health, National Center for Chronic Disease Prevention and Health Promotion, CDC.
- Pérez-Pérez, E., E. Serván-Mori, G. Nigenda, L. Ávila-Burgos, and D. Mayer-Foulkes (2019): “Government expenditure on health and maternal mortality in México: A spatial-econometric analysis,” *The International journal of health planning and management*, 34, 619–635.
- Pop-Eleches, C. (2010): “The Supply of Birth Control Methods, Education, and Fertility: Evidence from Romania,” *Journal of Human Resources*, 45, 971–997.
- Pourette, D., C. M. nd Rila Ratovoson, and P. Raharimalala (2018): “Complications with use of misoprostol for abortion in Madagascar: between ease of access and lack of information,” *Contraception*, 97, 116–121.
- Rambachan, A. and J. Roth (2019): “An Honest Approach to Parallel Trends,” mimeo, Harvard University.
- Rees, H., J. Katzenellenbogen, R. Shabodien, R. Jewkes, S. Fawcus, J. McIntyre, C. Lombard, and H. Truter (1997): “The epidemiology of incomplete abortion in South Africa,” *South African Medical Journal*, 87, 432–437.
- Rodríguez-Aguilar, R. (2018): “Maternal mortality in Mexico, beyond millennial development objectives: An age-period-cohort model,” *PloS one*, 13, e0194607.
- Sahatci, E. (1993): “Legal abortion improved women’s health in Albania.” *Newsletter (Women’s Global Network on Reproductive Rights)*, 10.
- Sánchez Fuentes, M. L., J. Paine, and B. Elliott-Buettner (2008): “The Decriminalisation of Abortion in Mexico City: How Did Abortion Rights Become a Political Priority?” *Gender and Development*, 16, 345–360.
- Schiavon, R., M. E. Collado, E. Troncoso, J. E. Soto Sánchez, G. O. Zorrilla, and T. Palermo (2010): “Characteristics of private abortion services in Mexico City after legalization,” *Reproductive health matters*, 18, 127–135.

- Schiavon, R., E. Troncoso, and G. Polo (2012a): “Analysis of maternal and abortion-related mortality in Mexico over the last two decades, 1990–2008,” *International Journal of Gynecology & Obstetrics*, 118, S78–S86.
- (2012b): “Analysis of maternal and abortion-related mortality in Mexico over the last two decades, 1990–2008,” *International Journal of Gynecology & Obstetrics*, 118, Supplement 2, S78 – S86.
- Serbanescu, F., L. Morris, P. Stupp, and A. Stanescu (1995): “The impact of recent policy changes on fertility, abortion, and contraceptive use in Romania,” *Studies in family planning*, 76–87.
- Seward, P. N., C. A. Ballard, and A. L. Ulene (1973): “The effect of legal abortion on the rate of septic abortion at a large county hospital,” *American Journal of Obstetrics & Gynecology*, 115, 335–338.
- Singh, K. and S. Ratnam (2015): “The influence of abortion legislation on maternal mortality,” *International Journal of Gynecology and Obstetrics*, 63, 123–129.
- Singh, S. and I. Maddow-Zimet (1999): “Facility-based treatment for medical complications resulting from unsafe pregnancy termination in the developing world, 2012: a review of evidence from 26 countries,” *BJOG: An International Journal of Obstetrics and Gynaecology*, 123, 1489–1498.
- Singh, S., L. Remez, and A. Tartaglione (2010): *Methodologies for estimating abortion incidence and abortion-related morbidity: a review*, Guttmacher Institute.
- Singh, S., G. Sedgh, A. Bankole, R. Hussain, and S. London (2012a): “Making abortion services accessible in the wake of legal reforms: A framework and six case studies,” Tech. rep., Guttmacher Institute.
- (2012b): “Making abortion services accessible in the wake of legal reforms: A framework and six case studies,” Tech. rep., Guttmacher Institute.
- Special Investigations Division, Committee of Government Reform, House of Representatives (2003): “Politics and Science in the Bush Administration. Prepared for Rep. Henry A. Waxman,” Available at: <http://web.archive.org/web/20121215150603/http://democrats.oversight.house.gov/images/stories/documents/20080130103545.pdf>. Accessed 2 June, 2018.
- Stephenson, P., M. Wagner, M. Badea, and F. Serbanescu (1992): “Commentary: the public health consequences of restricted induced abortion—lessons from Romania,” *American Journal of Public Health*, 82, 1328–1331.
- Stewart, G. K. and P. J. Goldstein (1971): “Effects on Septic Abortion and Maternal Mortality,” *Obstetrics & Gynecology*, 37, 510–514.
- Suh, S. (2014): “Rewriting abortion: Deploying medical records in jurisdictional negotiation over a forbidden practice in Senegal,” *Social Science & Medicine*, 108, 20–33.
- Sully, E., A. Biddlecom, J. Darroch, L. Ashford, N. Lince-Deroche, T. Riley, et al. (2019): “Adding It Up: Investing in Sexual and Reproductive Health,” *New York, NY: Guttmacher Institute*.
- Tuiran, R., V. Partida, O. Mojarro, and E. Zúñiga (2004): “Fertility in Mexico: trends and forecast,” in *Expert Group Meeting on Completing the Fertility Transition*, United Nations. Department of Economic and Social Affairs. Population Division. New York, New York, United Nations, Department of Economic and Social Affairs, Population Division, Compilation, 483–506.
- UNICEF (2005): “The ‘Rights’ Start to Life: A statistical analysis of birth registration,” Unicef publications, UNICEF.
- United Nations (2014): “Abortion Policies and Reproductive Health around the World 2014,” .

- Valente, C. (2014): “Access to abortion, investments in neonatal health, and sex-selection: Evidence from Nepal,” *Journal of Development Economics*, 107, 225–243.
- Velasco Sustaita, B. B. (Undated): “Recomendación de Códigos CIE-10a Diagnósticos Médicos en Escenarios a Gran Escala,” Thesis, Instituto Nacional de Astrofísica, Óptica y Electrónica.
- Venator, J. and J. Fletcher (2019): “Undue Burden Beyond Texas: An Analysis of Abortion Clinic Closures, Births, And Abortions in Wisconsin,” NBER Working Papers 26362, National Bureau of Economic Research, Inc.
- Winikoff, B. and W. R. Sheldon (2012): “Abortion: what is the problem?” *The Lancet*, 379, 594 – 596.
- World Health Organization (1987): “Studying Maternal Mortality in Developing Countries: A Guidebook,” Tech. Rep. HO/FHE/87.7, WHO.
- (2011): “Unsafe abortion: Global and regional estimates of the incidence of unsafe abortion and associated mortality in 2008,” Edition 6, WHO Library.
- (2018): “Preventing unsafe abortion,” Accessed at <http://www.who.int/news-room/fact-sheets/detail/preventing-unsafe-abortion> (consulted August 28, 2018).